

Neutral and cationic ruthenium hydrotris(pyrazolyl)borate derivatives containing bulky monodentate phosphines. Crystal structures of $[\text{RuTp}(\text{H}_2\text{O})(\text{PPr}^i_2\text{Me})_2][\text{CF}_3\text{SO}_3]\cdot\text{EtOH}$ and $[\text{RuTp}(\text{N}_2)(\text{PEt}_3)_2][\text{BPh}_4]$

Miguel A. Jiménez Tenorio, Manuel Jiménez Tenorio, M. Carmen Puerta* and Pedro Valerga

Departamento de Ciencia de Materiales e Ingeniería Metalúrgica y Química Inorgánica, Facultad de Ciencias, Universidad de Cádiz, Apartado 40, 11510 Puerto Real, Cádiz, Spain.
 E-mail: carmen.puerta@uca.es

Received 26th June 1998, Accepted 28th August 1998

The complexes $[\text{RuTp}(\text{Cl})(\text{PPr}^i_2\text{Me})_2]$ **1** and $[\text{RuTp}(\text{Cl})(\text{PEt}_3)_2]$ **2** were prepared by thermal displacement of PPh_3 from $[\text{RuTp}(\text{Cl})(\text{PPh}_3)_2]$ by the corresponding phosphine. A series of cationic complexes of the type $[\text{RuTp}(\text{L})(\text{PR}_3)_2]^+$ ($\text{L} = \text{H}_2\text{O}$, N_2 or CNBu^t ; $\text{PR}_3 = \text{PPr}^i_2\text{Me}$ or PEt_3) was obtained by chloride abstraction from **1** or **2** in the presence of L . As consequence of the steric crowding, one of the PPr^i_2Me ligands in **1** is labile, and it was readily replaced by neutral molecules such as MeCN or CNBu^t to yield the neutral complexes $[\text{RuTp}(\text{Cl})\text{L}(\text{PPr}^i_2\text{Me})]$ ($\text{L} = \text{MeCN}$ or CNBu^t). The monohydrides $[\text{RuTp}(\text{H})(\text{PPr}^i_2\text{Me})_2]$ and $[\text{RuTp}(\text{H})(\text{PEt}_3)_2]$ were obtained by treatment of **1** or **2** with NaBH_4 in MeOH . Protonation of these monohydrides led to the cationic dihydrogen complexes $[\text{RuTp}(\text{H}_2)(\text{PPr}^i_2\text{Me})_2]^+$ and $[\text{RuTp}(\text{H}_2)(\text{PEt}_3)_2]^+$ which were isolated as $[\text{BPh}_4]^-$ salts, and characterized by determination of their $(T_1)_{\text{min}}$ and $^1J_{\text{HD}}$ coupling constants in the corresponding isotopomers. The neutral hydride(dihydrogen) complex $[\text{RuTp}(\text{H})(\text{H}_2)(\text{PPr}^i_2\text{Me})]$, which resulted from the reaction of $[\text{RuTp}(\text{Cl})(\text{MeCN})(\text{PPr}^i_2\text{Me})]$ with NaBH_4 in MeOH , was characterized in analogous fashion. The crystal structures of the complexes $[\text{RuTp}(\text{H}_2\text{O})(\text{PPr}^i_2\text{Me})_2][\text{CF}_3\text{SO}_3]\cdot\text{EtOH}$ and $[\text{RuTp}(\text{N}_2)(\text{PEt}_3)_2][\text{BPh}_4]$ were determined.

Introduction

The chemistry of ruthenium hydrotris(pyrazolyl)borate (Tp) complexes, which had been rather underdeveloped compared to that of the related and formally homologous cyclopentadienyl and pentamethylcyclopentadienyl ruthenium derivatives, has attracted increasing attention in recent years. New TpRu complexes containing a variety of co-ligands, particularly nitrogen donors such as $\text{Me}_2\text{NCH}_2\text{CH}_2\text{NMe}_2$, MeCN , pyridine, *etc.* as well as cycloocta-1,5-diene (COD) and hemilabile phosphino amine ligands of the type $\text{Ph}_2\text{PCH}_2\text{CH}_2\text{NR}_2$, have been reported very recently,¹⁻⁶ and many of these have shown to be active catalysts for processes such as the dimerization of terminal acetylenes or the coupling of phenylacetylene with benzoic acid or allyl alcohols.^{1,7} Tertiary phosphines have also been used as co-ligands,⁸⁻¹⁵ and complexes such as $[\text{RuTp}(\text{X})(\text{PPh}_3)_2]$ ($\text{X} = \text{Cl}$ or H)^{8,12,15} and $[\text{RuTp}(\text{MeCN})(\text{PPh}_3)_2][\text{BF}_4]$ ¹² shown to be quite efficient catalyst precursors.^{7,12,16} Interestingly, complexes containing bidentate phosphine ligands, *e.g.* $[\text{RuTp}(\text{Cl})(\text{dppe})]$,¹ are catalytically inactive.⁷ It seems that the catalyst precursor must contain at least one weakly bound ligand which upon dissociation generates the co-ordinatively unsaturated, catalytically active species. We have previously prepared TpRu complexes containing the bulky diphosphine 1,2-bis(diisopropylphosphino)ethane (dippe), and studied its reactivity towards small molecules,¹⁵ and also towards terminal alkynes and alkynols.¹⁷ Although these complexes exhibit a chemical reactivity similar to that of their half-sandwich homologues,¹⁸ they are catalytically inactive. For this reason, and continuing our studies on the chemistry of ruthenium complexes with bulky phosphine ligands, we have now focused our attention on PPr^i_2Me , a bulky phosphine which can be considered as the monodentate equivalent of dippe. Dissociation of one monodentate PPr^i_2Me ligand is expected to occur more easily than the chelate ring opening in complexes containing

bidentate phosphines such as dippe or dppe, and hence such derivatives are more likely to act as suitable precursors for catalytically active species. In this work we describe the synthesis, properties and chemical reactivity of a range of neutral and cationic TpRu complexes containing PPr^i_2Me , including dinitrogen adducts as well as “classical” and “non-classical” hydrides, and also related derivatives with PEt_3 , a less sterically demanding phosphine ligand having electron-donating capabilities similar to those of PPr^i_2Me .

Experimental

All synthetic operations were performed under a dry dinitrogen or argon atmosphere following conventional Schlenk or dry-box techniques. Tetrahydrofuran, diethyl ether and light petroleum (boiling point range 40–60 °C) were distilled from the appropriate drying agents. All solvents were deoxygenated immediately before use. Triethylphosphine was purchased from Aldrich, whereas PPr^i_2Me was obtained by reaction of PClPr^i_2 (Aldrich) with MgMeI in diethyl ether; KTp^{19} and $[\text{RuTp}(\text{Cl})(\text{PPh}_3)_2]$ ⁸ were obtained according to published procedures. Infrared spectra were recorded in Nujol mulls on Perkin-Elmer FTIR Spectrum 1000 spectrophotometers, NMR spectra on Varian Unity 400 MHz or Gemini 200 MHz equipment. Chemical shifts are given in ppm from SiMe_4 (^1H and $^{13}\text{C}\{-^1\text{H}\}$) or 85% H_3PO_4 ($^{31}\text{P}\{-^1\text{H}\}$). The coupling constants $^3J_{\text{HH}}$ for the Tp ligand were all in the range 2–2.5 Hz. Microanalyses were by the Serveis Científico-Tècnics, Universitat de Barcelona.

Preparations

$[\text{RuTp}(\text{Cl})(\text{PPr}^i_2\text{Me})_2]$ **1.** To a solution of $[\text{RuTp}(\text{Cl})(\text{PPh}_3)_2]$ (1.75 g, 2 mmol) in toluene (15 ml), PPr^i_2Me (0.75 ml, *ca.* 5 mmol) was added *via* a syringe. The mixture was refluxed for 1 h, then allowed to cool to room temperature and light petrol-

eum (15 ml) added. A pale yellow, crystalline precipitate was obtained. It was filtered off, washed with diethyl ether and light petroleum, and dried *in vacuo*. Yield: 0.74 g, 60% (Found: C, 45.2; H, 7.28; N, 13.4. Calc. for $C_{23}H_{44}BClN_6P_2Ru$: C, 45.0; H, 7.17; N, 13.7%). IR: $\nu(\text{BH})$ 2456 cm^{-1} . NMR (C_6D_6): ^1H , δ (298 K) -0.23, 0.74, 1.34, 1.54 {m, $\text{P}[\text{CH}(\text{CH}_3)_2]_2$ }; 1.14 (d, $J_{\text{HP}} = 3.2$ Hz, PCH_3); 2.31, 2.95 {m, $\text{P}[\text{CH}(\text{CH}_3)_2]_2$ }; 5.85 (t, 1 H), 5.93 (t, 2 H), 7.29 (d, 1 H), 7.51 (d, 2 H), 7.54 (d, 1 H) and 8.39 (d, 2 H); δ (198 K, CD_2Cl_2) -0.63, 0.78, 0.92, 1.14, 1.25, 1.32, 1.39, 2.10, 3.35 (br, PPr^i_2Me); 6.05 (s, 1 H), 6.13 (br, 2 H), 7.14 (s, 1 H), 7.65 (br, 2 H), 7.70 (s, 1 H) and 7.91 (br, 2 H); ^{31}P - $\{^1\text{H}\}$, δ (298 K) 27.5 (s); δ (183 K, CD_2Cl_2) 23.9 (d), 25.9 (d), $^2J_{\text{PP}} = 30.9$ Hz; ^{13}C - $\{^1\text{H}\}$, δ (298 K) 6.8 (m, PCH_3); 16.2, 17.7, 19.1, 19.2 {s, $\text{P}[\text{CH}(\text{CH}_3)_2]_2$ }; 24.1, 28.1 {m, $\text{P}[\text{CH}(\text{CH}_3)_2]_2$ }; 104.7, 105.0, 134.5, 136.2, 145.1 and 147.1 [s, $\text{HB}(\text{C}_3\text{H}_3\text{N}_2)_3$].

[RuTp(Cl)(PEt₃)₂] 2. Complex 2 was prepared in a fashion analogous to that for 1, starting from [RuTp(Cl)(PPh₃)₂] (1.75 g, 2 mmol) and PEt₃ (0.75 ml, *ca.* 5 mmol). Yield: 0.68 g, 65% (Found: C, 43.0; H, 6.96; N, 14.1. Calc. for $C_{21}H_{40}BClN_6P_2Ru$: C, 43.1; H, 6.83; N, 14.4%). IR: $\nu(\text{BH})$ 2469 cm^{-1} . NMR (CDCl_3): ^1H , δ 0.75 [m, $\text{P}(\text{CH}_2\text{CH}_3)_3$], 1.87 [m, $\text{P}(\text{CH}_2\text{CH}_3)_3$]; 6.07 (t, 1 H), 6.13 (t, 2 H), 7.41 (d, 1 H), 7.61 (d, 2 H), 7.71 (d, 1 H) and 7.97 (d, 2 H); ^{31}P - $\{^1\text{H}\}$, δ 26.8 (s); ^{13}C - $\{^1\text{H}\}$, δ 7.8 [s, $\text{P}(\text{CH}_2\text{CH}_3)_3$], 18.4 [m, $\text{P}(\text{CH}_2\text{CH}_3)_3$]; 104.0, 105.2, 134.9, 136.2, 143.9 and 147.6 [s, $\text{HB}(\text{C}_3\text{H}_3\text{N}_2)_3$].

[RuTp(H₂O)(PPrⁱ₂Me)₂][CF₃SO₃] \cdot EtOH 3. To a tetrahydrofuran solution (15 ml) of complex 1 (0.3 g, 0.5 mmol) under argon, AgO_3SCF_3 (0.12 g, *ca.* 0.5 mmol) was added. The mixture was stirred at room temperature for 1 h, then, filtered through Celite in order to remove the precipitate of AgCl. The solvent was removed *in vacuo*, and the residue dissolved in 96% EtOH. Concentration and cooling to -20°C for several days afforded well formed crystals, which were filtered off, washed with light petroleum and dried *in vacuo*. Yield: 0.29 g, 77% (Found: C, 41.0; H, 6.83; N, 10.9. Calc. for $C_{26}H_{52}F_3N_6O_5P_2RuS$: C, 41.1; H, 6.85; N, 11.1%). IR: $\nu(\text{BH})$ 2463 cm^{-1} . NMR (CDCl_3): ^1H , δ -0.23, 1.01, 1.23, 1.34 {m, $\text{P}[\text{CH}(\text{CH}_3)_2]_2$ }; 1.45 (d, $J_{\text{HP}} = 6$ Hz, PCH_3); 2.26, 2.44 {m, $\text{P}[\text{CH}(\text{CH}_3)_2]_2$ }; 3.10 (br, H_2O); 6.08 (t, 1 H), 6.25 (t, 2 H), 7.29 (d, 1 H), 7.63 (d, 1 H), 7.73 (br, 2 H) and 7.84 (br, 2 H); ^{31}P - $\{^1\text{H}\}$, δ 28.5 (s); ^{13}C - $\{^1\text{H}\}$, δ 7.7 (m, PCH_3); 16.9, 18.7, 19.8, 20.0 {s, $\text{P}[\text{CH}(\text{CH}_3)_2]_2$ }; 25.1, 28.9 {m, $\text{P}[\text{CH}(\text{CH}_3)_2]_2$ }; 105.6, 106.0, 135.6, 137.3, 145.5 and 148.6 [s, $\text{HB}(\text{C}_3\text{H}_3\text{N}_2)_3$].

[RuTp(H₂O)(PEt₃)₂][BPh₄] 4. Complex 4 was obtained following a procedure identical to that for 3, starting from 2 (0.29 g, 0.5 mmol). It was converted into its tetraphenylborate salt by addition of NaBPh₄ (0.3 g, excess) to an ethanol solution. Cooling to -20°C afforded white crystals, which were filtered off, washed with light petroleum and dried *in vacuo*. Yield: 0.35 g, 80% (Found: C, 61.0; H, 7.14; N, 9.2. Calc. for $C_{45}H_{62}B_2N_6O_2P_2Ru$: C, 60.9; H, 6.99; N, 9.5%). IR: $\nu(\text{BH})$ 2479 cm^{-1} . NMR (CDCl_3): ^1H , δ 0.70 [m, $\text{P}(\text{CH}_2\text{CH}_3)_3$], 1.68 [m, $\text{P}(\text{CH}_2\text{CH}_3)_3$]; 5.30 (s, H_2O); 6.19 (t, 1 H), 6.24 (t, 2 H), 7.38 (d), 7.44 (s br), 7.69 (d, 2 H) and 7.77 (d, 1 H); ^{31}P - $\{^1\text{H}\}$, δ 25.0 (s); ^{13}C - $\{^1\text{H}\}$ [(CD_3)₂CO], δ 6.9 [s, $\text{P}(\text{CH}_2\text{CH}_3)_3$], 17.9 [m, $\text{P}(\text{CH}_2\text{CH}_3)_3$]; 106.0, 106.2, 136.0, 137.2, 143.2 and 147.9 [s, $\text{HB}(\text{C}_3\text{H}_3\text{N}_2)_3$].

[RuTp(N₂)(PPrⁱ₂Me)₂][BPh₄] 5. To a tetrahydrofuran solution (15 ml) of complex 1 (0.3 g, 0.5 mmol) under dinitrogen, AgO_3SCF_3 (0.12 g, *ca.* 0.5 mmol) was added. The mixture was stirred at room temperature for 2 h then filtered through Celite or centrifuged. The solvent was removed *in vacuo*, and the residue dissolved in MeOH. Addition of solid NaBPh₄ (0.3 g, excess), concentration and cooling to -20°C for several days afforded red crystals, which were filtered off, washed with ethanol and light petroleum and dried *in vacuo*. They were

recrystallized from a mixture of dichloromethane and light petroleum. Yield: 0.31 g, 68% (Found: C, 61.2; H, 7.03; N, 11.8. Calc. for $C_{47}H_{64}B_2N_8P_2Ru$: C, 61.0; H, 6.92; N, 12.1%). IR: $\nu(\text{BH})$ 2491, $\nu(\text{N}\equiv\text{N})$ 2159 cm^{-1} . NMR [(CD_3)₂CO]: ^1H , δ -0.32, 1.04, 1.32 {m, $\text{P}[\text{CH}(\text{CH}_3)_2]_2$ }; 1.68 (d, $J_{\text{HP}} = 7.2$ Hz, PCH_3); 2.36, 2.72 {m, $\text{P}[\text{CH}(\text{CH}_3)_2]_2$ }; 6.23 (t, 1 H), 6.33 (t, 2 H), 7.75 (d, 1 H), 7.84 (d, 2 H), 7.92 (d, 1 H) and 7.99 (d, 2 H); ^{31}P - $\{^1\text{H}\}$, δ 25.4 (s); ^{13}C - $\{^1\text{H}\}$, δ 6.9 (m, PCH_3); 16.5, 18.2, 18.8, 19.4 {s, $\text{P}[\text{CH}(\text{CH}_3)_2]_2$ }; 24.8, 27.0 {m, $\text{P}[\text{CH}(\text{CH}_3)_2]_2$ }; 106.9, 107.5, 137.3, 139.0, 146.5 and 150.7 [s, $\text{HB}(\text{C}_3\text{H}_3\text{N}_2)_3$].

[RuTp(N₂)(PEt₃)₂][BPh₄] 6. A procedure identical to that for the preparation and recrystallization of complex 5 was followed, starting from 2 (0.29 g, 0.5 mmol). Yield: 0.33 g, 73% (Found: C, 59.9; H, 7.01; N, 12.1. Calc. for $C_{45}H_{60}B_2N_8P_2Ru$: C, 60.2; H, 6.69; N, 12.5%). IR: $\nu(\text{BH})$ 2505 $\nu(\text{N}\equiv\text{N})$ 2163 cm^{-1} . NMR [(CD_3)₂CO]: ^1H , δ 0.78 (m, PCH_2CH_3), 1.93 (m, PCH_2CH_3); 6.20 (t, 1 H), 6.34 (t, 2 H), 7.70 (d, 1 H), 7.82 (d, 2 H), 7.91 (d, 1 H) and 7.98 (d, 2 H); ^{31}P - $\{^1\text{H}\}$, δ 25.8 (s); ^{13}C - $\{^1\text{H}\}$, δ (CD_2Cl_2) 7.9 [s, $\text{P}(\text{CH}_2\text{CH}_3)_3$], 18.5 [m, $\text{P}(\text{CH}_2\text{CH}_3)_3$]; 107.5, 106.5, 137.5, 138.8, 143.2 and 143.3 [s, $\text{HB}(\text{C}_3\text{H}_3\text{N}_2)_3$].

[RuTp(CNBU^t)(PPrⁱ₂Me)₂][BPh₄] 7. *Method A.* To a suspension of complex 1 (0.15 g, *ca.* 0.25 mmol) in EtOH (10 ml), NaBPh₄ (0.2 g, excess) and a few drops of CNBU^t were added. The mixture was heated smoothly for *ca.* 2 h using a water-bath. Then, the resulting yellow solution was concentrated and cooled to -20°C . The white solids were collected by filtration, washed with EtOH and light petroleum and dried *in vacuo*. Yield: 0.2 g, 82%.

Method B. A dichloromethane solution (10 ml) of complex 5 (0.1 g, *ca.* 0.11 mmol) was treated with a few drops of CNBU^t. The solution became pale yellow. It was stirred at room temperature for 10 min. Concentration and addition of light petroleum yielded a white precipitate, which was filtered off, washed with light petroleum and dried *in vacuo*. Yield: 0.1 g, quantitative (Found: C, 63.8; H, 7.55; N, 9.8. Calc. for $C_{52}H_{73}B_2N_7P_2Ru$: C, 63.7; H, 7.45; N, 10.0%). IR: $\nu(\text{BH})$ 2487, $\nu(\text{C}\equiv\text{N})$ 2124 cm^{-1} . NMR (CDCl_3): ^1H , δ -0.33, 0.91, 1.28, 1.36 {m, $\text{P}[\text{CH}(\text{CH}_3)_2]_2$ }; 1.29 (d, $J_{\text{HP}} = 6.8$ Hz, PCH_3); 1.44 [s, RuCNC(CH_3)₃]; 1.94, 2.41 {m, $\text{P}[\text{CH}(\text{CH}_3)_2]_2$ }; 6.20 (t, 2 H), 6.33 (t, 1 H), 7.63 (d, 2 H), 7.65 (d, 2 H), 7.69 (d, 1 H) and 7.83 (d, 1 H); ^{31}P - $\{^1\text{H}\}$, δ 26.0 (s); ^{13}C - $\{^1\text{H}\}$, δ 7.2 (m, PCH_3); 19.5, 19.1, 18.1, 16.2 {s, $\text{P}[\text{CH}(\text{CH}_3)_2]_2$ }; 30.8, 24.9 {m, $\text{P}[\text{CH}(\text{CH}_3)_2]_2$ }; 31.1 [s, RuCNC(CH_3)₃]; 49.3 [s, CNC(CH_3)₃]; 106.7, 136.8, 137.5, 144.6, 145.7 [s, $\text{HB}(\text{C}_3\text{H}_3\text{N}_2)_3$]; RuCNC(CH_3)₃ not observed.

[RuTp(CNBU^t)(PEt₃)₂][BPh₄] 8. This complex was obtained by either of the two procedures outlined for the preparation of 7, starting either from 2 (method A) or from 6 (method B), with similar yields (Found: C, 63.1; H, 7.33; N, 10.1. Calc. for $C_{50}H_{69}B_2N_7P_2Ru$: C, 63.1; H, 7.25; N, 10.3%). IR: $\nu(\text{BH})$ 2471, $\nu(\text{C}\equiv\text{N})$ 2123 cm^{-1} . NMR (CDCl_3): ^1H , δ 0.72 [m, $\text{P}(\text{CH}_2\text{CH}_3)_3$]; 1.43 [s, CNC(CH_3)₃]; 1.69 [m, $\text{P}(\text{CH}_2\text{CH}_3)_3$]; 6.02 (t, 2 H), 6.31 (t, 1 H), 7.48 (d, 2 H), 7.62 (d, 1 H), 7.66 (d, 2 H) and 7.84 (d, 1 H); ^{31}P - $\{^1\text{H}\}$, δ 24.0 (s); ^{13}C - $\{^1\text{H}\}$, δ 7.5 [s, $\text{P}(\text{CH}_2\text{CH}_3)_3$], 19.1 [m, $\text{P}(\text{CH}_2\text{CH}_3)_3$]; 30.8 [s, RuCNC(CH_3)₃]; 49.5 [s, RuCNC(CH_3)₃]; 106.4, 106.5, 136.5, 137.1, 143.4 and 145.5 [s, $\text{HB}(\text{C}_3\text{H}_3\text{N}_2)_3$]; RuCNC(CH_3)₃ not observed.

[RuTp(Cl)(MeCN)(PPrⁱ₂Me)] 9. A solution of complex 1 (0.3 g, 0.5 mmol) in MeCN (15 ml) was heated under reflux for 2 h. Then, the solvent was removed *in vacuo*. The resulting yellow microcrystalline material was washed with several portions of light petroleum and dried thoroughly. Yield: 0.24 g, 95% (Found: C, 41.2; H, 5.89; N, 18.5. Calc. for $C_{18}H_{30}BClN_7PRu$: C, 41.4; H, 5.74; N, 18.8%). IR: $\nu(\text{BH})$ 2479; $\nu(\text{C}\equiv\text{N})$ 2269, 2245 cm^{-1} . NMR [(CD_3)₂CO]: ^1H , δ 0.15, 1.10, 1.35 {m, $\text{P}[\text{CH}(\text{CH}_3)_2]_2$ }; 1.52 (d, $J_{\text{HP}} = 8$ Hz, PCH_3); 2.24 {m, $\text{P}[\text{CH}(\text{CH}_3)_2]_2$ }; 2.56 (s, RuNCCH₃); 6.11 (t, 1 H), 6.16 (t, 1 H), 6.17 (t, 1 H),

7.56 (d, 1 H), 7.68 (d, 1 H), 7.69 (d, 1 H), 7.74 (d, 1 H), 7.80 (d, 1 H) and 7.88 (d, 1 H); ^{31}P - $\{^1\text{H}\}$, δ 42.1 (s); ^{13}C - $\{^1\text{H}\}$, δ 4.1 (s, RuNCCH₃); 4.2 (d, $J_{\text{CP}} = 21.7$, PCH₃); 16.5 (d, $J_{\text{HP}} = 1.3$), 16.7 (d, $J_{\text{HP}} = 1.3$), 17.8, 17.9 {s, P[CH(CH₃)₂]₂}; 24.7 {d, $J_{\text{CP}} = 22.7$ Hz, P[CH(CH₃)₂]₂}; 105.6, 105.7, 106.6, 134.4, 136.2, 136.5, 142.1, 146.0 and 146.2 [s, HB(C₃H₃N₂)₃].

[RuTp(Cl)(CNBu^t)(PPrⁱ₂Me)] 10. To a solution of complex **1** (0.15 g, ca. 0.25 mmol) in tetrahydrofuran a few drops of CNBu^t were added. The mixture was heated at 60 °C for 2 h. Then the solvent was removed *in vacuo*, leaving a yellow powder which was washed with several portions of light petroleum and dried *in vacuo*. Yield: 0.14 g, quantitative (Found: C, 44.3; H, 6.40; N, 17.1. Calc. for C₂₁H₃₆BClN₇PRu: C, 44.7; H, 6.38; N, 17.4%). IR: $\nu(\text{BH})$ 2460, $\nu(\text{C}\equiv\text{N})$ 2071, 2099 cm⁻¹. NMR [(CD₃)₂CO]: ^1H , δ 0.31, 1.00, 1.04, 1.14 {m, P[CH(CH₃)₂]₂}; 0.85 (d, $J_{\text{HP}} = 3.2$ Hz, PCH₃); 1.51 [s, CNC(CH₃)₃]; 1.99, 2.19 {m, P[CH(CH₃)₂]₂}; 6.15 (t, 1 H), 6.16 (t, 1 H), 6.24 (t, 1 H), 7.50 (d, 1 H), 7.65 (d, 1 H), 7.79 (d, 1 H), 7.82 (d, 1 H), 7.84 (br, 1 H) and 7.94 (d, 1 H); ^{31}P - $\{^1\text{H}\}$, δ 42.5 (s); ^{13}C - $\{^1\text{H}\}$ (C₆D₆), δ 4.2 (d, $J_{\text{CP}} = 40$, PCH₃); 15.9, 17.1, 17.7, 19.2 {s, P[CH(CH₃)₂]₂}; 24.1 (d, $J_{\text{CP}} = 41$), 28.5 {d, $J_{\text{CP}} = 49$ Hz, P[CH(CH₃)₂]₂}; 31.0 [s, RuCNC(CH₃)]; 48.3 [s, RuCNC(CH₃)₃]; 105.0, 105.1, 105.5, 134.8, 135.5, 142.6, 143.9, 145.3 [s, HB(C₃H₃N₂)₃]; 233 [br, RuCNC(CH₃)₃].

[RuTp(Cl)(PPrⁱ₂Me)(PEt₃)] 11. To a tetrahydrofuran solution of complex **1** (0.15 g, 0.25 mmol) neat PEt₃ (0.05 ml, slight excess) was added. The mixture was heated at 60 °C for 1 h. Removal of the solvent and washing with light petroleum afforded a yellow solid, which was dried *in vacuo*. Yield: 0.13 g, quantitative (Found: C, 43.9; H, 7.15; N, 14.0. Calc. for C₂₂H₄₂BClN₆P₂Ru: C, 44.1; H, 7.01; N, 14.0%). IR: $\nu(\text{BH})$ 2473 cm⁻¹. NMR (CDCl₃): ^1H , δ 0.78 [m, P(CH₂CH₃)₃]; 1.88 [m, P(CH₂CH₃)₃]; -0.39, 0.97, 1.39, 1.47 {m, P[CH(CH₃)₂]₂}; 1.37 (d, $J_{\text{HP}} = 7.2$ Hz, PCH₃); 2.71, 2.26 {m, P[CH(CH₃)₂]₂}; 6.08 (t, 1 H), 6.13 (br, 2 H), 7.44 (d, 1 H), 7.61 (d, 1 H), 7.71 (br, 2 H), 8.00 (d, 2 H) and 8.03 (d, 2 H); ^{31}P - $\{^1\text{H}\}$, δ 31.1 (d); 23.3 (d), $^2J_{\text{PP}} = 32.3$ Hz; ^{13}C - $\{^1\text{H}\}$, δ 6.8 (d, $J_{\text{CP}} = 34.8$, PCH₃); 6.9 [s, P(CH₂CH₃)₃]; 18.5 [d, $J_{\text{CP}} = 45.4$, P(CH₂CH₃)₃]; 19.3, 19.2, 17.8, 15.7 {s, P[CH(CH₃)₂]₂}; 24.0, 28.9 {d, $J_{\text{CP}} = 42.4$ Hz, P[CH(CH₃)₂]₂}; 104.0, 105.1, 105.3, 134.8, 134.9, 136.4, 144.0, 144.4 and 147.4 [s, HB(C₃H₃N₂)₃].

[RuTp(H)(PPrⁱ₂Me)₂] 12. To a slurry of complex **1** (0.3 g, 0.5 mmol) in MeOH (15 ml) an excess of NaBH₄ (0.15 g) was added. The mixture was heated using a water-bath, until effervescence ceased. After 45 min a yellow-orange solution was obtained. Removal of the solvent yielded an oily residue, which was dissolved in light petroleum and filtered through Celite. Concentration and cooling to -20 °C afforded a yellow solid which was filtered off and dried *in vacuo*. This compound slowly turns green on standing under dinitrogen or argon, even when stored in a freezer. Yield: 0.16 g, 58% (Found: C, 47.3; H, 7.92; N, 14.1. Calc. for C₂₃H₄₅BN₆P₂Ru: C, 47.7; H, 7.77; N, 14.5%). IR: $\nu(\text{BH})$ 2456, $\nu(\text{RuH})$ 1917 cm⁻¹. NMR (C₆D₆): ^1H , δ -15.36 (t, $^2J_{\text{HP}} = 28.8$, RuH); 0.02, 0.64, 1.15, 1.24 {m, P[CH(CH₃)₂]₂}; 1.05 (d, $J_{\text{HP}} = 5.8$ Hz, PCH₃); 2.16 {m, P[CH(CH₃)₂]₂}; 5.70 (s br, 2 H), 5.99 (s br, 1 H), 7.00 (s br, 2 H), 7.37 (s br, 1 H), 7.59 (s br, 1 H) and 7.76 (s br, 2 H); ^{31}P - $\{^1\text{H}\}$, δ 47.8 (s); ^{13}C - $\{^1\text{H}\}$, δ 9.2 (m, PCH₃); 17.8, 18.1, 18.2, 19.3 {s, P[CH(CH₃)₂]₂}; 25.3, 31.9 {m, P[CH(CH₃)₂]₂}; 104.5, 104.6, 135.0, 145.4 and 146.9 [s, HB(C₃H₃N₂)₃].

[RuTp(H)(PEt₃)₂] 13. Complex **13** was obtained in a fashion analogous to that for **12**, starting from **2** (0.29 g, 0.5 mmol). Yield: 0.16 g, 60% (Found: C, 45.4; H, 7.27; N, 14.9. Calc. for C₂₁H₄₁BN₆P₂Ru: C, 45.8; H, 7.44; N, 15.3%). IR: $\nu(\text{BH})$ 2456, $\nu(\text{RuH})$ 1915 cm⁻¹. NMR (C₆D₆): ^1H , δ -15.11 (t, $^2J_{\text{HP}} = 29.2$ Hz, RuH); 0.74 [m, P(CH₂CH₃)₃]; 1.50 [m, P(CH₂CH₃)₃]; 5.86

(t, 2 H), 6.14 (t, 1 H), 7.52 (d, 2 H), 7.75 (s br, 3H) and 8.25 (d, 1 H); ^{31}P - $\{^1\text{H}\}$, δ 45.9 (s); ^{13}C - $\{^1\text{H}\}$, δ 8.1 [s, P(CH₂CH₃)₃]; 21.2 [m, P(CH₂CH₃)₃]; 105.0, 105.8, 135.2, 145.8 and 146.9 [s, HB(C₃H₃N₂)₃].

[RuTp(H₂)(PPrⁱ₂Me)₂][BPh₄] 14. To a solution of complex **12** (0.15 g, 0.26 mmol) in diethyl ether (10 ml) at -80 °C under argon, HBF₄·OEt₂ (2–3 drops, excess) was added. The mixture was allowed to warm to room temperature. Then the solvent was removed *in vacuo*, and the residue treated with a MeOH solution containing NaBPh₄ (0.2 g, excess). A white, microcrystalline precipitate was obtained. It was filtered off, washed with ethanol and light petroleum and dried *in vacuo*. This compound was recrystallized from a dichloromethane–ethanol mixture under argon. The isotopomer [RuTp(HD)(PPrⁱ₂Me)]⁺ was generated *in situ*, by reaction of **12** with DBF₄·OEt₂ (obtained from D₂O–HBF₄·OEt₂ 3:1). Yield: 0.19 g, 81% (Found: C, 62.5; H, 7.46; N, 9.1. Calc. for C₄₇H₆₆B₂N₆P₂Ru: C, 62.8; H, 7.34; N, 9.35%). IR: $\nu(\text{BH})$ 2489 cm⁻¹. NMR (CD₂Cl₂): ^1H , δ -9.55 [s br, Ru(H₂); (T_1)_{min} 16 ms at 203 K, 400 MHz, $^1J_{\text{HD}} = 31.1$, $J_{\text{HP}} = 7.3$ Hz]; -0.08, 0.90, 1.35 {m, P[CH(CH₃)₂]₂}; 1.42 (d, PCH₃); 2.00, 2.22 {m, P[CH(CH₃)₂]₂}; 6.23 (t, 2 H), 6.45 (t, 1 H), 7.68 (d, 2 H), 7.72 (d, 2 H), 7.77 (d, 1 H) and 7.96 (d, 1 H); ^{31}P - $\{^1\text{H}\}$, δ 30.6 (s); ^{13}C - $\{^1\text{H}\}$, δ 7.0 (m, PCH₃); 16.6, 17.6, 18.5, 19.4 {s, P[CH(CH₃)₂]₂}; 25.1, 30.3 {m, P[CH(CH₃)₂]₂}; 107.0, 107.5, 137.4, 138.2, 146.7 and 147.0 [s, HB(C₃H₃N₂)₃].

[RuTp(H₂)(PEt₃)₂][BPh₄] 15. A procedure identical to that for complex **14** was followed for **15**, starting from **13** (0.15 g, 0.27 mmol). Yield: 0.17 g, 75% (Found: C, 61.9; H, 7.18; N, 9.4. Calc. for C₄₅H₆₂B₂N₆P₂Ru: C, 62.0; H, 7.12; N, 9.65%). IR: $\nu(\text{BH})$ 2495 cm⁻¹. NMR (CD₂Cl₂): ^1H , δ -9.83 [s br, Ru(H₂), (T_1)_{min} 18 ms at 203 K, 400 MHz, $^1J_{\text{HD}} = 30.9$, $J_{\text{HP}} = 7.1$ Hz]; 0.78 [m, P(CH₂CH₃)₃]; 1.74 [m, P(CH₂CH₃)₃]; 6.23 (t, 2 H), 6.41 (t, 1 H), 7.59 (d, 2 H), 7.71 (d, 2 H), 7.78 (d, 1 H) and 7.93 (d, 1 H); ^{31}P - $\{^1\text{H}\}$, δ 28.1 (s); ^{13}C - $\{^1\text{H}\}$, δ 7.3 [s, P(CH₂CH₃)₃]; 19.4 [m, P(CH₂CH₃)₃]; 107.0, 107.2, 146.9, 146.2, 137.2 and 138.0 [s, HB(C₃H₃N₂)₃].

[RuTp(H₃)(PPrⁱ₂Me)] 16. A solution of complex **9** (0.15 g, 0.29 mmol) in MeOH (10 ml) was treated with an excess of NaBH₄ (0.15 g). The mixture was heated at 60 °C for 1 h. Then the solvent was removed *in vacuo*, the residue extracted with light petroleum, and the solution filtered through Celite. A sticky residue was obtained upon solvent removal and dissolved in warm MeOH. Filtration, concentration and cooling to -20 °C afforded pale yellow crystals, which were filtered off and dried *in vacuo*. A mixture of the isotopomers [RuTp(H₂D)(PPrⁱ₂Me)] and [RuTp(H)D₂(PPrⁱ₂Me)] was obtained by gentle heating of a CD₃OD solution of **16**. Yield: 0.1 g, 77% (Found: C, 42.3; H, 6.91; N, 18.5. Calc. for C₁₆H₃₀BN₆PRu: C, 42.8; H, 6.68; N, 18.7%). IR: $\nu(\text{BH})$ 2475, $\nu(\text{RuH})$ 1946 cm⁻¹. NMR (C₆D₆): ^1H , δ -10.45 [d, $^2J_{\text{HP}} = 19.6$, RuH₃, (T_1)_{min} 41 ms at 201 K, CD₂Cl₂, 400 MHz, $^1J_{\text{HD}} = 7.4$]; 0.69, 0.83 {m, P[CH(CH₃)₂]₂}; 1.03 (d, $J_{\text{HP}} = 7.2$ Hz, PCH₃); 5.99 (s br), 7.56 (s br) and 7.84 (s br); ^{31}P - $\{^1\text{H}\}$, δ 62.9 (s); ^{13}C - $\{^1\text{H}\}$, δ 13.3 (d, $J_{\text{CP}} = 44$, PCH₃); 15.0, 18.3, 18.4, 18.5 {s, P[CH(CH₃)₂]₂}; 22.6, 26.5 {d, $J_{\text{CP}} = 44$ Hz, P[CH(CH₃)₂]₂}; 105.7, 135.6 and 146.2 [s br, HB(C₃H₃N₂)₃].

Crystallography

Crystals suitable for X-ray diffraction analysis were mounted onto a glass fiber and transferred to an AFC6S-Rigaku automatic diffractometer ($T = 290$ K, Mo-K α radiation, graphite monochromator, $\lambda = 0.71073$ Å). Accurate unit cell parameters and an orientation matrix in each case were determined by least-squares fitting from the settings of 25 high-angle reflections. Crystal data and details on data collection and refinements are given in Table 1. Data were collected by the ω -2 θ scan method in both cases. Lorentz-polarization corrections

were applied. Decay was monitored by measuring three standard reflections every 100 measurements. Decay and semiempirical absorption correction (ψ method) were also applied. The structures were solved by Patterson methods and subsequent expansion of the models using DIRDIF.²⁰ Reflections having $I > 3\sigma(I)$ were used for structure refinement. For complex **3** all non-hydrogen atoms were anisotropically refined; H(1), H(2) and H(52) were localized in Fourier-difference maps and the remaining hydrogen atoms included at idealized positions and not refined. In the case of compound **6** all the non-hydrogen atoms in the cation except the phosphine ethyl groups were anisotropically refined, and the remaining non-hydrogen atoms were isotropically refined. Hydrogen atoms were included at idealized positions and not refined. Since the space groups were non-centrosymmetrical in both cases, the two possible enantiomorphs were checked and no significant differences found. All calculations for data reduction, structure solution, and refinement were carried out on a VAX 3520 computer at the Servicio Central de Ciencia y Tecnología de la Universidad de Cádiz, using the TEXSAN²¹ software system and ORTEP²² for plotting. Maximum and minimum peaks in the final Fourier-difference maps were $+1.42$ and -0.97 e \AA^{-3} for **3**, and $+0.56$ and -0.43 e \AA^{-3} for **6**.

CCDC reference number 186/1139.

See <http://www.rsc.org/suppdata/dt/1998/3601/> for crystallographic files in .cif format.

Results and discussion

The complexes $[\text{RuTp}(\text{Cl})(\text{PPr}^i_2\text{Me})_2]$ **1** and $[\text{RuTp}(\text{Cl})(\text{PEt}_3)_2]$ **2** were obtained by thermal displacement of PPh_3 from $[\text{RuTp}(\text{Cl})(\text{PPh}_3)_2]$ by the corresponding phosphine in refluxing toluene, a procedure which has been previously used for the synthesis of $[\text{RuTp}(\text{Cl})(\text{dippe})]$.¹⁵ The physical properties of these pale yellow, crystalline compounds match those of the dippe derivative. The presence of six separate pyrazole ring proton resonances in the ^1H NMR spectra, and of one singlet in the $^{31}\text{P}\{-^1\text{H}\}$ NMR spectra, of these complexes at room temperature suggest an octahedral structure analogous to that found for the parent complex $[\text{RuTp}(\text{Cl})(\text{PPh}_3)_2]$ by X-ray crystallography. It is interesting that the complexes $[\text{Ru}(\text{C}_5\text{R}_5)\text{Cl}(\text{PEt}_3)_2]$ ($\text{R} = \text{H}$ or Me), formal homologues of **2**, are known. However, attempts made to synthesize organometallic counterparts of complex **1**, namely $[\text{Ru}(\text{C}_5\text{R}_5)\text{Cl}(\text{PPr}^i_2\text{Me})_2]$ ($\text{R} = \text{H}$ or Me), have been unsuccessful.²³ It seems that the bulk of PPr^i_2Me makes difficult, or even prevents, the formation of half-sandwich species containing two of these phosphine ligands in a cisoid disposition, an observation which is consistent with the fact that complexes of the type $[\text{Ru}(\text{C}_5\text{R}_5)\text{Cl}(\text{PR}_3)_2]$ ($\text{R} = \text{H}$ or Me ; $\text{PR}_3 = \text{PPh}_3$ or PCy_3) are also unknown.²⁴ Instead, 16-electron species $[(\text{C}_5\text{Me}_5)\text{RuCl}(\text{PR}_3)]$ ($\text{PR}_3 = \text{PCy}_3$, PPr^i_3 , PPhPr^i_2) have shown to be stable, yet reactive.²⁵ Given the increased bulk of the Tp ligand compared to C_5H_5 or C_5Me_5 , the formation of a complex containing two *cis*- PPr^i_2Me ligands such as **1** is remarkable. Compound **1** shows in its ^1H NMR spectrum one multiplet at $\delta -0.23$ attributable to CH_3 protons of isopropyl groups of the PPr^i_2Me ligand, which is somewhat unusual, since such a high-field resonance has not been observed in the ^1H NMR spectra of other TpRu phosphine complexes, e.g. **2** or $[\text{RuTp}(\text{Cl})(\text{dippe})]$. This anomalous chemical shift for isopropyl groups casts some doubts on whether **1** is really a plain six-co-ordinate species having a κ^3 -Tp ligand (A), or a highly fluxional five-co-ordinate molecule containing a κ^2 -Tp group and phosphines in a *trans* disposition, being stabilized by means of an "agostic" interaction with the isopropyl group of one of the phosphine ligands (B).

Ruthenium derivatives containing one κ^2 -Tp ligand and two *trans* phosphine ligands have recently been reported, i.e. $[\text{Ru}(\kappa^2\text{-Tp})\text{H}(\text{CO})(\text{PPr}^i_3)_2]$ ²⁶ and $[\text{Ru}(\kappa^2\text{-Tp})\text{H}(\text{CO})(\text{PPh}_3)_2]$,²⁷ so this is a real possibility to be considered in our case. When the





Fig. 1 The NMR spectra of complex **1** at 183 K (CD_2Cl_2): (A) $^{31}\text{P}\{-^1\text{H}\}$, (B) ^1H . The peak marked with * corresponds to the solvent.



temperature is lowered the $^{31}\text{P}\{-^1\text{H}\}$ NMR resonance observed in the spectrum of **1** broadens, and splits into two separate signals which resolve into doublets at 183 K [Fig. 1(A)], suggesting the non-equivalence of the phosphorus atoms at this temperature. At this temperature, three of the pyrazole ring resonances in the ^1H NMR spectrum broaden [Fig. 1(B)], although they do not split as would be expected if the slow exchange limit were reached to give rise to nine separate signals, indicative of the non-equivalence of the three pyrazole rings of the Tp ligand. Apart from broadening and shifting to $\delta -0.63$, lowering the temperature causes no effect on the high-field phosphine proton resonance. Apparently, this spectral behaviour might be consistent with a formulation as a κ^2 -Tp complex with "agostic" interaction. However, no reduced $^1J_{\text{CH}}$ coupling constants were observed in the proton-coupled ^{13}C NMR spectrum, as would be expected if "agostic" hydrogen atoms were present. If instead we assume that **1** is just a sterically crowded six-co-ordinate complex, the unusual chemical shift observed might be caused by anisotropy due to the fact that some of the isopropyl hydrogen atoms of the phosphine, as result of the steric pressure, are forced near the magnetic ring current influence of a pyrazole ring. The inequivalent phosphines at low temperature may arise from freezing out conformers around single M-P and P-C bonds. No crystals of **1** suitable for X-ray diffraction could be obtained. However, crystals of the aqua complex $[\text{RuTp}(\text{H}_2\text{O})(\text{PPr}^i_2\text{Me})_2][\text{CF}_3\text{SO}_3]\cdot\text{EtOH}$ **3** were serendipitously obtained during an attempt to prepare the co-ordinatively unsaturated complex $[\text{RuTp}(\text{PPr}^i_2\text{Me})_2][\text{CF}_3\text{SO}_3]$ by reaction of **1** with AgO_3SCF_3 in EtOH, and its structure was determined.

A structural ORTEP view of complex **3** is shown in Fig. 2. Selected bond lengths and angles are listed in Table 2. Com-


Table 1 Summary of data for the crystal structure analysis of complexes **3** and **6**

	3	6
Formula	C ₂₆ H ₅₂ N ₆ BF ₃ O ₅ P ₂ RuS	C ₄₅ H ₆₀ B ₂ N ₈ P ₂ Ru
<i>M</i>	791.62	897.66
Crystal size/mm	0.32 × 0.25 × 0.14	0.40 × 0.36 × 0.22
Crystal system	Monoclinic	Orthorhombic
Space group	<i>P</i> 2 ₁ / <i>c</i> (no. 14)	<i>P</i> 2 ₁ 2 ₁ 2 ₁ (no. 19)
<i>a</i> /Å	14.337(5)	15.908(4)
<i>b</i> /Å	22.077(6)	11.550(3)
<i>c</i> /Å	13.204(6)	22.555(4)
β /°	90.44(2)	
<i>U</i> /Å ³	3734(1)	4637(2)
<i>Z</i>	4	4
<i>D</i> _c /g cm ⁻³	1.408	1.286
μ (Mo-K α)/cm ⁻¹	6.03	4.37
<i>F</i> (000)	1648	1880
Unique reflections	6012	3754
Observed reflections (<i>I</i> > 3 σ _{<i>i</i>})	3434	1998
Number of parameters	406	338
<i>R</i>	0.062	0.066
<i>R</i> ' (<i>w</i> = σ_F^{-2})	0.075	0.077

Table 2 Selected bond distances (Å) and angles (°) for [RuTp(H₂O)(PPrⁱ₂Me)₂][CF₃SO₃] \cdot EtOH

Ru(1)–P(1)	2.342(3)	O(1)⋯O(5)	2.59(1)
Ru(1)–P(2)	2.362(3)	O(1)–H(1)	1.21
Ru(1)–O(1)	2.142(6)	O(1)–H(2)	0.95
Ru(1)–N(12)	2.049(7)	O(5)–H(3)	0.97
Ru(1)–N(22)	2.147(8)	O(1)–H(3)	1.70
Ru(1)–N(32)	2.140(7)	O(2)–H(1)	1.83
O(1)⋯O(2)	2.82(1)		
P(1)–Ru(1)–P(2)	98.54(10)	O(1)–Ru(1)–N(12)	169.7(3)
P(1)–Ru(1)–O(1)	92.3(2)	O(1)–Ru(1)–N(22)	85.1(3)
P(1)–Ru(1)–N(12)	92.6(2)	O(1)–Ru(1)–N(32)	83.6(3)
P(1)–Ru(1)–N(22)	170.8(2)	N(12)–Ru(1)–N(22)	88.6(3)
P(1)–Ru(1)–N(32)	91.5(2)	N(12)–Ru(1)–N(32)	87.3(3)
P(2)–Ru(1)–O(1)	96.1(2)	N(22)–Ru(1)–N(32)	79.4(3)
P(2)–Ru(1)–N(12)	92.1(2)	Ru(1)–O(1)–H(1)	120.6
P(2)–Ru(1)–N(22)	90.5(2)	Ru(1)–O(1)–H(2)	110.0
P(2)–Ru(1)–N(32)	170.0(2)	H(1)–O(1)–H(2)	88.4

compound **3**, which also shows a high field phosphine proton resonance at δ –0.23, has a six-coordinate distorted octahedral structure, with phosphines in a cisoid arrangement showing no evidence for “agostic” interaction with the metal. The water molecule appears bound to ruthenium, and linked to an oxygen atom of the [CF₃SO₃][–] anion *via* a hydrogen bond. The OH group of the ethanol solvate also forms a hydrogen bond with the co-ordinated water molecule. The hydrogen bond distances O⋯O are 2.59(1) and 2.82(1) Å. Strong hydrogen bonds between water and the [CF₃SO₃][–] anion have also been observed in the recently reported crystal structures of [RuTp(H₂O){Ph₂P(CH₂)₂NMe₂}[CF₃SO₃] \cdot 0.5CH₂Cl₂³ and [RuTp(H₂O)(COD)][CF₃SO₃]² which consist of neutral dimeric units linked by hydrogen bonds. In these compounds the hydrogen bond O⋯O distances range from 2.714 Å to 2.966 Å, comparing well with the O(1)⋯O(2) separation, although the O(1)⋯O(5) bond distance of 2.59(1) Å is indicative of a much stronger hydrogen bond between the water ligand and the ethanol solvate molecule. The Ru(1)–O(1) bond distance 2.142(6) Å is similar to the Ru–O separations observed for [RuTp(H₂O){Ph₂P(CH₂)₂NMe₂}[CF₃SO₃] \cdot 0.5CH₂Cl₂ [2.190(2) Å]³ and [RuTp(H₂O)(COD)][CF₃SO₃] [2.151(4) Å]² and also for the complex [Ru{HC(pz)₃}(H₂O)₃][*p*-MeC₆H₄SO₃] \cdot 1.5 H₂O [2.131(1) Å; HC(pz)₃ = tris(pyrazolyl)methane].²⁸ The angle P(1)–Ru(1)–P(2) of 98.54(10)° is significantly larger than the values of *ca.* 85° found for complexes containing the bidentate phosphine dippe, such as [RuTp{C(OMe)CH₂CO₂Me}(dippe)][BPh₄]¹⁷ or [RuTp(H₂)(dippe)][BPh₄]¹⁵ since the phos-


**Fig. 2** An ORTEP drawing of the compound [RuTp(H₂O)(PPrⁱ₂Me)₂][CF₃SO₃] \cdot EtOH with 50% probability thermal ellipsoids. Hydrogen atoms, except those of the water ligand, are omitted.

phine ligands are monodentate in **3** and do not have the “bite angle” imposed by the backbone ethane chain in dippe. In this fashion the steric repulsions between the two PPrⁱ₂Me are minimized, but this forces some of the methyl groups on the isopropyl substituents [C(11) and C(3)] to move towards the gap between two pyrazole rings. As consequence, the hydrogen atoms attached to these methyl groups fall into the the magnetic ring current influence of the pyrazole rings, giving rise to the anomalous high field chemical shift observed for these protons in the ¹H NMR spectra. From the crystal structure of **3** it is clear that two bulky PPrⁱ₂Me can appear simultaneously bound to a TpRu centre, and therefore the absence of known complexes of the type [Ru(C₅Me₅)Cl(PR₃)₂] (PR₃ = bulky phosphine ligand: PPrⁱ₃, PCy₃, PPrⁱ₂Me, PPrⁱ₂Ph *etc.*) might be due to electronic rather than to steric reasons.^{24,25}

The aqua complex [RuTp(H₂O)(PET₃)₂][BPh₄]⁴ was also obtained by chloride abstraction from **2** under argon in the presence of water, using AgO₃SCF₃. The water protons in **3** and **4** appear respectively at δ 3.1 and 5.30 respectively. If the chloride abstraction from complexes **1** and **2** is performed under dinitrogen instead of argon, then the dinitrogen adducts [RuTp(N₂)(PPrⁱ₂Me)₂]⁺ and [RuTp(N₂)(PET₃)₂]⁺ are obtained, which were isolated as their respective tetraphenylborate salts **5** and **6**. These compounds display one strong ν (N₂) band at 2159 and 2163 cm⁻¹ respectively. Since our initial report of the synthesis of the dinitrogen complex [RuTp(N₂)(dippe)][BPh₄][ν (N₂) 2165 cm⁻¹],¹⁵ two other TpRu dinitrogen adducts have been described: [RuTp(N₂){Ph₂P(CH₂)₂NMe₂}[CF₃SO₃][ν (N₂) 2182 cm⁻¹]³ and [RuTp(N₂)(PPh₃)₂][BF₄][ν (N₂) 2177 cm⁻¹],¹² which suggest that, at variance with their cyclopentadienyl or pentamethylcyclopentadienyl homologues,^{18a} TpRu dinitrogen complexes seem to be a well established class of compounds. The crystal structure of **6** was determined. An ORTEP view of the complex cation is shown in Fig. 3. Selected bond lengths and angles are listed in Table 3. As for compound **3**, the coordination around the Ru atom is distorted octahedral. The dinitrogen ligand is bound in the end-on manner, with a Ru(1)–N(1)–N(2) angle of 166(3)°. The Ru(1)–N(1) and N(1)–N(2) separations are 1.91(2) and 1.01(2) Å respectively, which are fully consistent with the dimensions obtained for other ruthenium dinitrogen complexes including [RuTp(N₂){Ph₂P(CH₂)₂NMe₂}[CF₃SO₃] (Ru–N 1.943(4), N–N 1.097(5) Å).³ As

Table 3 Selected bond distances (Å) and angles (°) for [RuTp(N₂)-(PEt₃)₂][BPh₄]

Ru(1)–P(1)	2.362(5)	Ru(1)–N(22)	2.15(2)
Ru(1)–P(2)	2.365(6)	Ru(1)–N(32)	2.07(1)
Ru(1)–N(1)	1.91(2)	N(1)–N(2)	1.01(2)
Ru(1)–N(12)	2.17(1)		
P(1)–Ru(1)–P(2)	99.8(2)	P(2)–Ru(1)–N(32)	88.9(5)
P(1)–Ru(1)–N(1)	91.8(7)	N(1)–Ru(1)–N(12)	90(1)
P(1)–Ru(1)–N(12)	90.7(5)	N(1)–Ru(1)–N(22)	87.5(9)
P(1)–Ru(1)–N(22)	172.7(5)	N(1)–Ru(1)–N(32)	176.5(9)
P(1)–Ru(1)–N(32)	91.6(5)	N(12)–Ru(1)–N(22)	82.0(7)
P(2)–Ru(1)–N(1)	91.2(9)	N(12)–Ru(1)–N(32)	88.7(7)
P(2)–Ru(1)–N(12)	169.3(5)	N(22)–Ru(1)–N(32)	89.0(7)
P(2)–Ru(1)–N(22)	87.5(5)	Ru(1)–N(1)–N(2)	166(3)

**Fig. 3** An ORTEP drawing of the cation [RuTp(N₂)(PEt₃)₂]⁺ with 50% probability thermal ellipsoids. Hydrogen atoms are omitted.

in most other cases, the observed N(1)–N(2) bond distance is identical within the experimental error to that of the free N₂ molecule, a fact which in the case of [RuTp(N₂){Ph₂P(CH₂)₂NMe₂}]⁺ has been interpreted in terms of a delicate compensatory influence of σ -bond strengthening and π -bond weakening in the N₂ molecule, as inferred from extended Hückel molecular orbital (EHMO) calculations.³ This also accounts for the high frequency at which the ν (N₂) IR band appears in these complexes. The dinitrogen ligand in **5** and **6** is labile, and easily replaceable by good neutral donors such as CNBu^t, leading to the cationic complexes [RuTp(CNBu^t)(PPrⁱ₂Me)₂][BPh₄]**7** [ν (CN) 2124 cm⁻¹] and [RuTp(CNBu^t)(PEt₃)₂][BPh₄]**8** [ν (CN) 2123 cm⁻¹], which have octahedral structures as inferred from spectral data and do not require further comment.

As a consequence of its bulkiness, PPrⁱ₂Me in complex **1** is substitutionally labile, at variance with PEt₃ in **2**. Thus, **1** reacts smoothly with neutral donors such as MeCN or CNBu^t furnishing the neutral complexes [RuTp(Cl)L(PPrⁱ₂Me)] (L = MeCN **9** or CNBu^t **10**). These compounds display strong bands in their respective IR spectra at 2269 and 2245 cm⁻¹, and at 2071 and 2099 cm⁻¹, attributable to ν (CN) in the ligands MeCN and CNBu^t. The three pyrazole rings of the Tp ligand in these species are inequivalent, and hence nine separate proton and carbon resonances appear in their ¹H and ¹³C-¹H NMR spectra. The characteristic high-field phosphine proton resonance observed for compounds of the type [RuTp(X)(PPrⁱ₂Me)₂] is absent in the ¹H NMR spectra of **9** and **10**. The steric crowding around the metal decreases upon replacement of one of the


PPrⁱ₂Me by a less bulky ligand, so the methyl protons of the isopropyl substituents are not forced any more to remain under the magnetic ring current influence of the pyrazole rings. Triethylphosphine also displaces one PPrⁱ₂Me from **1**, affording the mixed phosphine derivative [RuTp(Cl)(PPrⁱ₂Me)(PEt₃)]**11**, which is characterized by the presence of two doublets in its ³¹P-¹H NMR spectrum corresponding to an AB spin system, as expected. In this particular compound the substitution of one PPrⁱ₂Me by one PEt₃ does not relieve the steric pressure and the high field Prⁱ proton resonance is still observed in the ¹H NMR spectrum. Other compounds of the type [RuTp(Cl)L(PR₃)] (L = MeCN, py, CO, P(OMe)₃ or PMe₃; PR₃ = PPh₃ or PCy₃) have been described recently, these being prepared either by displacement of DMF from [RuTp(Cl)(DMF)(PPh₃)] by L,¹ or by reaction of L with the ruthenium(III) complex [RuTp(Cl)(OR)(PCy₃)] (R = Me or Et).⁹

Synthesis and characterization of hydride complexes

Complex **1** and **2** reacted with NaBH₄ in MeOH affording the neutral monohydride complexes [RuTp(H)(PPrⁱ₂Me)₂]**12** and [RuTp(H)(PEt₃)₂]**13**. These air-sensitive compounds display one strong ν (RuH) IR band near 1915 cm⁻¹, and one triplet in their ¹H NMR spectra attributable to the hydride proton. As result of the smaller size of hydrogen compared to chloride or other atoms, there is less steric pressure in the hydride complexes containing two PPrⁱ₂Me ligands than in other [RuTp(X)(PPrⁱ₂Me)₂] derivatives, and therefore all phosphine proton resonances in the ¹H NMR spectrum of **12** have positive chemical shifts relative to tetramethylsilane.

Both complexes **12** and **13** are protonated by HBF₄·OEt₂ furnishing the dihydrogen adducts [RuTp(H₂)(PPrⁱ₂Me)₂]⁺ and [RuTp(H₂)(PEt₃)₂]⁺ which were isolated as their tetraphenylborate salts **14** and **15**. The dihydrogen ligand in these complexes is characterized by a broad resonance in the corresponding ¹H NMR spectra, having short minimum longitudinal relaxation times ($T_{1\text{min}}$) of 16 and 18 ms respectively (400 MHz). The coupling constants ¹J_{HD} observed for the isotopomers [RuTp(HD)(PPrⁱ₂Me)₂]⁺ and [RuTp(HD)(PEt₃)₂]⁺ are 31.1 and 30.9 Hz. These values of ($T_{1\text{min}}$) and ¹J_{HD} compare well with those previously found for other cationic TpRu dihydrogen complexes such as [RuTp(H₂)(dippe)][BPh₄]¹⁵ [RuTp(H₂)(PPh₃)₂][BF₄]¹² and also [RuTp(H₂)(CO)(PPrⁱ₃)]-[BF₄]²⁶ being consistent with the presence of a co-ordinated dihydrogen molecule. From the values of ($T_{1\text{min}}$), a separation d (H–H) of 0.95 Å for both **14** and **15** has been estimated, assuming fast spinning of the dihydrogen ligand. Complexes **14** and **15** are white, crystalline solids, indefinitely stable at room temperature under argon. Under dinitrogen the dihydrogen ligand is displaced very slowly by N₂ to yield the corresponding dinitrogen complex **5** or **6**. As for other cationic TpRu derivatives, equilibrium or irreversible tautomerization to the ruthenium(IV) dihydride form has not been observed, this being attributed to the particular electron donating capabilities of the Tp group (*e.g.* in comparison with those of the formally related C₅H₅ or C₅Me₅ ligands), as well as to the fact that this ligand favours six- over seven-co-ordinate species.^{12,15,26}

We attempted to prepare the neutral monohydride complex [RuTp(H)(MeCN)(PPrⁱ₂Me)] by treatment of **9** with NaBH₄ in MeOH. However, in the course of the reaction the MeCN ligand was lost, and the ultimate product obtained turned out to be the hydride(dihydrogen) complex [RuTp(H)(H₂)(PPrⁱ₂Me)]**16**, which was isolated in the form of pale yellow crystals. This compound displays one strong IR band at 1946 cm⁻¹ attributable to ν (RuH). The ¹H NMR spectrum shows one doublet at δ –10.45 (²J_{HHP} = 19.6 Hz, 3 H) corresponding to the hydridic protons. No decoalescence of this resonance is observed as the temperature is lowered, suggesting that there is rapid atom exchange between hydride and dihydrogen sites, leading to the equivalence of these protons.



Only three rather broad proton and carbon pyrazole ring resonances are observed for complex **16** even at low temperature, indicative of the equivalence of the three pyrazole rings of the Tp ligand due to fluxional behaviour. Accordingly, the $^{31}\text{P}\{-^1\text{H}\}$ NMR spectrum shows one singlet. The value of $(T_1)_{\text{min}}$ for the hydride resonance is 40.8 ms (208 K, 400 MHz). As consequence of rapid chemical exchange with the terminal hydride, this relaxation time is averaged, but in agreement with the presence of one $\eta^2\text{-H}_2$ ligand within the complex. Complex **16** undergoes H–D exchange with CD_3OD , leading to the isotopomers $[\text{RuTp}(\text{H})(\text{HD})(\text{PPr}_2\text{Me})]$ and $[\text{RuTp}(\text{D})(\text{HD})(\text{PPr}_2\text{Me})]$. A $^1J_{\text{HD}}$ coupling constant of 7.4 Hz is observed, which is also averaged. Assuming a rapid hydride–dihydrogen fluxionality in a $\text{MH}(\text{H}_2)$ system, $^1J_{\text{HD}}$ of the dihydrogen ligand is equal to 3 times the observed $^1J_{\text{HD}}$ coupling constant,¹⁴ and hence in our case turns out to be 22.2 Hz, a value which is typical for dihydrogen co-ordinated to a transition metal. The values of $(T_1)_{\text{min}}$ and $^1J_{\text{HD}}$ for **16** are very similar to those found for the compounds $[\text{RuTp}(\text{H})(\text{H}_2)(\text{PR}_3)]$ ($\text{PR}_3 = \text{PCy}_3$ ¹³ or PPh_3 ²⁹). The derivatives $[\text{RuTp}^*(\text{H})(\text{H}_2)(\text{PCy}_3)]$ [$\text{Tp}^* = \text{tris}(3,5\text{-dimethylpyrazolyl})\text{hydroborate}$ or $\text{tris}(4\text{-bromo-3-isopropylpyrazolyl})\text{hydroborate}$] have also been described,¹⁴ and their spectral properties match those of **16**. It is interesting that the compounds $[\text{RuTp}^*(\text{H})(\text{H}_2)(\text{PCy}_3)]$ ¹⁴ and $[\text{RuTp}(\text{H})(\text{H}_2)(\text{PPh}_3)]$ ²⁹ have been obtained by hydrogenation of suitable precursor complexes, using pressures of H_2 ranging from ca. 3 ($\text{PR}_3 = \text{PCy}_3$) to 6–40 atm ($\text{PR}_3 = \text{PPh}_3$). In our case the hydride–dihydrogen complex is formed smoothly just by reaction of **9** with NaBH_4 in MeOH, the use of a H_2 atmosphere not being required. Under similar conditions, but with longer reaction times, the reaction of $[\text{RuTp}(\text{Cl})(\text{MeCN})(\text{PPh}_3)]$ with NaBH_4 in MeOH leads to the hydrido carbonyl complex $[\text{RuTp}(\text{H})(\text{CO})(\text{PPh}_3)]$. In general, the reaction of $[\text{RuTp}(\text{H})(\text{MeCN})(\text{PPh}_3)]$ with NaBH_4 and alcohols RCH_2OH ($\text{R} = \text{Me}$, Et , Ph or tolyl) yields alkyl or aryl TpRu carbonyl derivatives $[\text{RuTp}(\text{R})(\text{CO})(\text{PPh}_3)]$ resulting from the decarbonylation of the alcohol.²⁹ However, we have not detected similar species so far in our system.

The chemical reactivity of hydride complexes **12–16** is currently being investigated in detail. In a forthcoming paper we will describe stoichiometric and catalytic C–C coupling reactions in 1-alkynes mediated by the complexes described in this work.

Acknowledgements

We wish to thank the Ministerio de Educación y Ciencia of Spain (Dirección General de Investigación Científica y Técnica, Project PB94 -1306) and Junta de Andalucía (PAI-FQM 0188) for financial support.

References

- C. Gemel, G. Trimmel, C. Slugovc, S. Kremel, K. Mereiter, R. Schmid and K. Kirchner, *Organometallics*, 1996, **15**, 3998.
- C. Gemel, P. Wiede, K. Mereiter, V. N. Sapunov, R. Schmid and K. Kirchner, *J. Chem. Soc., Dalton Trans.*, 1996, 4071.
- G. Trimmel, C. Slugovc, P. Wiede, K. Mereiter, V. N. Sapunov, R. Schmid and K. Kirchner, *Inorg. Chem.*, 1997, **36**, 1076.
- C. Slugovc, P. Wiede, K. Mereiter, R. Schmid and K. Kirchner, *Organometallics*, 1997, **16**, 2768.
- K. Kirchner, personal communication.
- F. A. Jalón, A. Otero and A. Rodríguez, *J. Chem. Soc., Dalton Trans.*, 1995, 1629.
- C. Slugovc, D. Doberer, C. Gemel, R. Schmid, K. Kirchner, B. Winkler and F. Stelzer, *Monatsh. Chem.*, 1998, **129**, 221.
- N. W. Alcock, I. D. Burns, K. S. Claire and A. F. Hill, *Inorg. Chem.*, 1992, **31**, 2906.
- C. Gemel, G. Kickelbick, R. Schmid and K. Kirchner, *J. Chem. Soc., Dalton Trans.*, 1997, 2113.
- C. Slugovc, V. Sapunov, P. Wiede, K. Mereiter, R. Schmid and K. Kirchner, *J. Chem. Soc., Dalton Trans.*, 1997, 4209.
- N.-Y. Sun and S. J. Simpson, *J. Organomet. Chem.*, 1992, **434**, 341.
- W.-C. Chan, C.-P. Lau, Y.-Z. Chen, Y.-Q. Fang, S.-M. Ng and G. Jia, *Organometallics*, 1997, **16**, 34.
- M. A. Halcrow, B. Chaudret and S. Trofimenko, *J. Chem. Soc., Chem. Commun.*, 1993, 465.
- B. Moreno, S. Sabo-Etienne, B. Chaudret, A. Rodríguez-Fernández, F. A. Jalón and S. Trofimenko, *J. Am. Chem. Soc.*, 1995, **117**, 7441.
- M. Jiménez-Tenorio, M. A. Jiménez Tenorio, M. C. Puerta and P. Valerga, *Inorg. Chim. Acta*, 1997, **259**, 77.
- C. Slugovc, K. Mereiter, E. Zobetz, R. Schmid and K. Kirchner, *Organometallics*, 1996, **15**, 5275.
- M. Jiménez-Tenorio, M. A. Jiménez Tenorio, M. C. Puerta and P. Valerga, *Organometallics*, 1997, **16**, 5528.
- (a) I. de los Ríos, M. Jiménez Tenorio, J. Padilla, M. C. Puerta and P. Valerga, *Organometallics*, 1996, **15**, 4565; (b) I. de los Ríos, M. Jiménez Tenorio, M. C. Puerta and P. Valerga, *J. Am. Chem. Soc.*, 1997, **119**, 6529 and refs. therein.
- S. Trofimenko, *Inorg. Synth.*, 1970, **12**, 99.
- P. T. Beurskens, DIRDIF, Technical Report 1984/1, Crystallography Laboratory, Toernooiveld, 1984.
- TEXSAN, Single-Crystal Structure Analysis Software, version 5.0, Molecular Structure Corporation, Houston, TX, 1989.
- C. K. Johnson, ORTEP, A Thermal Ellipsoid Plotting Program, Oak Ridge National Laboratory, Oak Ridge, TN, 1965.
- M. Jiménez Tenorio, J. Padilla, M. C. Puerta and P. Valerga, unpublished work.
- B. K. Campion, R. H. Heyn and T. D. Tilley, *J. Chem. Soc., Chem. Commun.*, 1988, 278; T. Arliguie, C. Border, B. Chaudret, J. Devillers and R. Poilblanc, *Organometallics*, 1989, **8**, 1308.
- T. J. Johnson, K. Foltig, W. E. Streib, J. D. Martin, J. C. Huffman, S. A. Jackson, O. Eisenstein and K. G. Caulton, *Inorg. Chem.*, 1995, **34**, 488.
- C. Bohanna, M. A. Esteruelas, A. V. Gómez, A. M. López and M. P. Martínez, *Organometallics*, 1997, **16**, 4464.
- I. D. Burns, A. F. Hill, A. J. P. White, D. J. Williams and J. D. E. T. Wilton-Ely, *Organometallics*, 1998, **17**, 1552.
- A. Llobet, D. J. Hodgson and T. J. Meyer, *Inorg. Chem.*, 1990, **29**, 3760.
- Y.-Z. Cheng, W.-C. Chan, C.-P. Lau, H.-S. Chu, H.-L. Lee and G. Jia, *Organometallics*, 1997, **16**, 1241.