

Nouvelle Cuisine for the Computation of the Annihilating Ideal of f^s

J. Gago-Vargas¹, M.I. Hartillo-Hermoso², and J.M. Ucha-Enríquez^{1,*}

¹ Depto. de Álgebra, Universidad de Sevilla. Apdo. 1160, E-41080 Sevilla, Spain
gago@algebra.us.es, ucha@us.es

² Depto. de Matemáticas, Universidad de Cádiz. Apdo. 40,
E-11510 Puerto Real, Spain
isabel.hartillo@uca.es

Abstract. Let f_1, \dots, f_p be polynomials in $\mathbf{C}[x_1, \dots, x_n]$ and let $D = D_n$ be the n -th Weyl algebra. The annihilating ideal of $f^s = f_1^{s_1} \cdots f_p^{s_p}$ in $D[s] = D[s_1, \dots, s_p]$ is a necessary step for the computation of the Bernstein-Sato ideals of f_1, \dots, f_p .

We point out experimental differences among the efficiency of the available methods to obtain this annihilating ideal and provide some upper bounds for the complexity of its computation.

1 Introduction

Fix two integers $n \geq 1, p \geq 1$ and two sets of variables (x_1, \dots, x_n) and (s_1, \dots, s_p) . Let us consider $f_1, \dots, f_p \in \mathbf{C}[x] = \mathbf{C}[x_1, \dots, x_n]$ and let $D = D_n$ be the n -th Weyl algebra. A polynomial $b(s) \in \mathbf{C}[s] = \mathbf{C}[s_1, \dots, s_p]$ is said to be a *Bernstein-Sato polynomial associated to f* if the following functional equation holds for a certain $P(s) \in D[s]$:

$$b(s)f^s = P(s)f^{s+1},$$

where $\mathbf{1} = (1, \dots, 1)$. These polynomials form an ideal called the *Bernstein-Sato ideal* \mathcal{B}_f , or simply \mathcal{B} to abbreviate. Analogous functional equations with respect to vectors different to $\mathbf{1}$ yield other different Bernstein-Sato ideals (see for example [Ba1]).

In [L1] it is proved that \mathcal{B} is not zero. This fact is a generalization of the classical proof of Bernstein ([Be1]) for the case $p = 1$, in which the generator of \mathcal{B} is called *the* Bernstein-Sato polynomial, $b_f(s)$. The analytical work was made in [Bj1] for $p = 1$ and in [Sa1],[Sa2] for $p > 1$.

The roots of $b_f(s)$ encode important algebro-geometrical data (see [Ma1], [H1] or [BS1] to mention only a few) and a complete understanding of all roots for a general f is open. For the case $p > 1$ there is a lot of work to do yet: there are conjectures on the primary decomposition of \mathcal{B} , on the conditions over f for \mathcal{B} to be principal, to be zero-dimensional, etc.

In [O1] was presented the first algorithm to find the Bernstein-Sato polynomial, and alternative methods have been proposed to obtain \mathcal{B} in the general case

* All authors partially supported by MTM2004-01165 and FQM-333.

in [OT1], [Ba1] and [BM1]. All these methods have a feature in common: their first step is the computation of the *annihilating ideal* of f^s in $D[s]$, $Ann_{D[s]}f^s$. We recall here some experimental evidences in favor of the method of Briançon-Maisonobe [BM1] with respect to the method of Oaku-Takayama [OT1].

Then we will give upper bounds of the complexity of computing $Ann_{D[s]}f^s$, the previous requirement for both algorithms. To obtain this bounds we use —as far as possible— the techniques and results of [Gr1] on the complexity of solving systems of linear equations over rings of differential operators (that extend the classical polynomial case treated in [Se1]). In particular, we show that the construction of Grigoriev can not be directly generalized to any non-commutative algebra, including the algebra proposed by Briançon-Maisonobe. We prove that the complexity of computing $Ann_{D[s]}f^s$ using the method of [BM1] is that of the calculation of a Gröbner basis in the n -th Weyl algebra with some extra p commutative variables ($2n + p$ variables at most), while in the case of the method [OT1] is the calculation of such a basis in a $(n + p)$ -th Weyl algebra with some extra $2p$ variables (so $2n + 4p$ variables in all).

We are very grateful to the referees for helping us to clarify our initial version.

2 Preliminaries

In this section we just remind briefly some details of the methods of Briançon-Maisonobe and Oaku-Takayama, respectively.

2.1 Method of Briançon-Maisonobe

In this case the computations are made in the non-commutative algebra

$$R = D_n[s, t] = D_n[s_1, \dots, s_p, t_1, \dots, t_p],$$

an extension of the n -th Weyl algebra D in which the new variables s, t satisfy the relations $[s_i, t_j] = \delta_{ij}t_i$. It is a *Poincaré-Birkhoff-Witt (PBW) algebra*:

Definition 1. *A PBW algebra R over a ring k is an associative algebra generated by finitely many elements x_1, \dots, x_n subject to the relations*

$$Q = \{x_j x_i = q_{ji} x_i x_j + p_{ji}, 1 \leq i < j \leq n\},$$

where each p_{ji} is a finite k -linear combination of standard terms $\mathbf{x}^\alpha = x_1^{\alpha_1} \dots x_n^{\alpha_n}$ and each $q_{ji} \in k$ with the two following conditions:

1. There is an admissible¹ order \prec on \mathbf{N}^n such that $\exp(p_{ji}) \prec \exp(x_j x_i)$ for every $1 \leq i < j \leq n$.
2. The standard terms \mathbf{x}^α , with $\alpha \in \mathbf{N}^n$, form a k -basis of R as a vector space.

¹ Here admissible means a total order among the elements of \mathbf{N}^n with $\mathbf{0}$ as least element.

It is possible to compute Gröbner bases in PBW algebras. The book [BGV1] is a good introduction to the subject of effective calculus in this fairly general family.

The following algorithm computes \mathcal{B} , starting from

$$I := \text{Ann}_R(f^s) = \langle s_j + f_j t_j, \partial_i + \sum_j \frac{\partial f_j}{\partial x_i} t_j, 1 \leq i \leq n, 1 \leq j \leq p \rangle.$$

Algorithm 1. You have to:

1. Obtain $J = \text{Ann}_{D_n[s]} f^s = \langle G_1 \cap D_n[s] \rangle$ where G_1 is a Gröbner basis of I with respect to any term ordering with variables t_j greater than the others (that is, an *elimination ordering for the t_j* .)
2. $\mathcal{B} = \langle G_2 \cap \mathbf{C}[s] \rangle$, where G_2 is a Gröbner basis of $J + (f_1, \dots, f_p)$ with respect to any term ordering with x_i, ∂_j greater than the s_l .

2.2 Method of Oaku-Takayama

All the computations are made in Weyl algebras. More precisely, starting from

$$I' = \langle t_j - f_j, \sum_{j=1}^p \frac{\partial f_j}{\partial x_i} \partial_{t_j} + \partial_i, i = 1, \dots, n, j = 1, \dots, p \rangle$$

Algorithm 2. You have to:

1. Obtain $J' = I' \cap \mathbf{C}[t_1 \partial_{t_1}, \dots, t_n \partial_{t_n}] \langle x, \partial_x \rangle$.
2. $J = \text{Ann}_{D_n[s]}(f^s) = J''$, where J'' denotes the ideal generated by the generators of J' after replacing each $t_i \partial_{t_i}$ by $-s_i - 1$.
3. $\mathcal{B} = \langle G_2 \cap \mathbf{C}[s] \rangle$, where G_2 is a Gröbner basis of $J + (f_1, \dots, f_p)$ with respect to any term ordering with x_i, ∂_j greater than the s_l .

The second step above² is again the elimination of all the variables but (s_1, \dots, s_p) . The computation of

$$I' \cap \mathbf{C}[t_1 \partial_{t_1}, \dots, t_n \partial_{t_n}] \langle x, \partial_x \rangle$$

uses $2n + 4p$ variables, as new variables u_j, v_j for $1 \leq j \leq p$ are introduced. More precisely, the first calculation is an elimination of these new variables for the ideal

$$\langle t_j - u_j f_j, \sum_{j=1}^p \frac{\partial f_j}{\partial x_i} u_j \partial_{t_j} + \partial_i, 1 - u_j v_j, 1 \leq i \leq n, 1 \leq j \leq p \rangle,$$

and some more technical steps must be followed (see [OT1, Procedure 4.1.]).

² Often the bottleneck to obtain the Bernstein-Sato ideal is this step. As far as we know, the example for $p = 2$ with $f_1 = x^2 + y^3, f_2 = x^3 + y^2$ is intractable for the available systems.

3 Experimental Data

Here we give some examples for the cases $p = 1, 2$ and $p > 2$ for which it is clear the superiority of Briançon-Maisonobe's method. They have been tested³ using SINGULAR::PLURAL 2.1 (see [GLS1]) in a PC Pentium IV, 1Gb RAM and 3.06GHz running under Windows XP.

SINGULAR::PLURAL 2.1 is a system for non-commutative general purpose, so the calculations in our algebras are not supposed to be optimal. We present the following data only for the sake of comparing both methods in the same system. In the case of [BM1] method we have used a pure lexicographical ordering, while for [OT1] we have used typical elimination ordering. These are the orderings with best results for each case.

The typical input for SINGULAR::PLURAL 2.1 looks like this for [BM1] method:

```
ring r = 0, (t(1..3), s(1..3), x, y, z, Dx, Dy, Dz), lp;

matrix C[12][12]=0;

C[1,4]=t(1);C[2,5]=t(2);C[3,6]=t(3);C[7,10]=1;C[8,11]=1;C[9,12]=1;

system("PLURAL",1,C);

poly f1 = x*z+y ; poly f2 = x*y+z; poly f3 = y*z+x;

ideal i =s(1)+t(1)*f1,s(2)+t(2)*f2,s(3)+t(3)*f3, Dx +
t(1)*diff(f1,x)+t(2)*diff(f2,x)+t(3)*diff(f3,x), Dy +
t(1)*diff(f1,y)+t(2)*diff(f2,y)+t(3)*diff(f3,y), Dz +
t(1)*diff(f1,z)+t(2)*diff(f2,z)+t(3)*diff(f3,z);

ideal I = std(i);
```

And this one for [OT1] method:

```
ring r = 0, (u, v, x, y, z, t, Dx, Dy, Dz, Dt), (a(1,1), dp);

matrix C[10][10]=0;

C[3,7]=1;C[4,8]=1;C[5,9]=1;C[6,10]=1;

system("PLURAL",1,C);
```

...

³ The CPU times must be considered as approximations: as it is explained in the SINGULAR::PLURAL 2.1 Manual, the command `timer` is not absolutely reliable due to the shortcomings of the Windows operating system.

- Case $p = 1$:** In the following examples $f \in \mathbf{C}[x, y]$ or $f \in \mathbf{C}[x, y, z]$. They have been chosen taking into account Arnold’s classification of singularities. **E** means the memory was exhausted and the system reported an error.

Table 1. CPU times for the computation of $\text{Ann}f^s$

f	Briançon-Maisonobe’s method	Oaku-Takayama’s method
$x^3 + xy^2 + z^2$	< 0.01s	0.39s
$x^4 + y^3 + z^2$	< 0.01s	0.39s
$yx^3 + y^3 + z^2$	0.06s	3.97s
$x^3 + y^2 + z^2$	< 0.01s	0.02s
$x^5 + y^2 + z^2$	< 0.01s	4.66s
$x^7 + y^2 + z^2$	< 0.01s	298.56s
$x^4 + y^5 + xy^4$	0.56s	E (> 12h)

- Case $p = 2$:** In Table 2 the examples f_1, f_2 are in $\mathbf{C}[x, y]$ or $\mathbf{C}[x, y, z]$.

Table 2. CPU times for the computation of $\text{Ann}f_1^{s_1} f_2^{s_2}$

f_1	f_2	Briançon-Maisonobe’s method	Oaku-Takayama’s method
$x^3 + y^2$	$x^2 + y^3$	0.72s	6363.97s
$x^5 + y^3$	$x^3 + y^5$	3.53s	E (> 6h)
$x^7 + y^5$	$x^5 + y^7$	11.84s	E (> 6h)
$x^3 + y^2$	$xz + y$	< 0.01s	9.73s
$x^5 + y^2$	$xz + y$	< 0.01s	1568.59 s
$x^{11} + y^5$	$xz + y$	3s	E (> 6h)

- Case $p > 2$:** In Table 3 we have some examples for more than two functions.

Table 3. CPU times for the computation of $\text{Ann}f_1^{s_1} \dots f_p^{s_p}$

f_1	f_2	f_3	Briançon-Maisonobe’s method	Oaku-Takayama’s method
$x + y$	$x - y$	$x^2 + y$	< 0.01s	29.46s
$x + y$	$x^2 + y$	$x + y^2$	2.64s	E
$x + y$	$x^2 + y$	$x^2 + y^3$	116.24s	E
$x + y$	$x^2 + y$	$x^3 + y^2$	1728.41s	E

When the single functions f_1, \dots, f_p are “simple” enough (for example, linear) it is possible to obtain $\text{Ann}_{D[s_1, \dots, s_p]} f_1^{s_1} \dots f_p^{s_p}$ for p rather big (say 15 or 20). This ideal can be related to the annihilating ideal of $f = f_1 \dots f_p$. This idea has been exploited with success in [GHU1] to compute annihilating ideals for f , where f defines very hard examples of arrangements of hyperplanes of theoretical interest. In Table 4 we compare the results of applying this idea in SINGULAR::PLURAL 2.1 with obtaining directly $\text{Ann}_{D[s]} f^s$ using the powerful system Asir (see [N1]).

Table 4. Some arrangements of hyperplanes

$f = f_1 \cdots f_p$	Briançon-Maisonobe's method computing $\text{Ann}_{D[s_1, \dots, s_p]} f_1^{s_1} \cdots f_p^{s_p}$	Asir computing $\text{Ann}_{D[s]} f^s$
$xyz(x+y)(x-y)(x-2y-z)$	0.62s	0.93s
$xyz(x-y)(x+y)(x-2y)$	0.05s	0.03s
$xyz(x+y)(x-y)(x+y-z)$	0.06s	3.54s
$xyzu(x+y+z+u)$	0.01s	6.99s
$xyzuv(x+y+z+u+v)$	0.02s	169L31s
$xyzuvw(x+y+z+u+v+w)$	0.05s	> 3 days

4 Complexity

In [Gr1] a bound for the degree of the solutions of a general system of linear equations over the Weyl algebra is given, with a procedure somewhat similar to the one of the commutative case of [Se1]. In this section we study how far the work of Grigoriev is applicable to our PBW algebra R of 2.1. His construction has two different steps: in the first, the given system is reduced to another system in a diagonal form. In the second, it is shown how to normalize the new system in order to eliminate, successively, the variables.

4.1 Diagonalization

We need three technicals lemma to reduce the system to a diagonal form. They generalize analogous lemmas of Grigoriev's paper (see [Gr1, Lemma 1]) and their proofs are, more or less, straightforward. Here deg means the *total degree* of a term, that is, the sum of the exponents of all its variables.

Lemma 1. *Let A be a $(m-1) \times m$ matrix with entries in a Poincaré-Birkhoff-Witt algebra S with a basis of p elements. If $\text{deg}(a_{ij}) \leq d$, there exists a nonzero-vector $f = (f_1, \dots, f_m) \in S^m$ such that $Af = 0$ and $\text{deg}(f) \leq 2p(m-1)d = N$.*

If we work in a noetherian domain (eventually non-commutative), we can always define the *rank* of a finite module as in [St1]. Given a square matrix in a Poincaré-Birkhoff-Witt algebra we say that it is *non-singular* if it has maximal rank, and in this case we can obtain a left quasi-inverse with the precedent lemma:

Lemma 2. *Given a $m \times m$ matrix B over a PBW algebra S as in Lemma 1, non-singular, it has a left quasi-inverse matrix G over S , such that $\text{deg}(G) \leq N$.*

Lemma 3. *Given a system of linear equations over a PBW algebra S , it is defined by a $m \times s$ matrix A of rank r , with its elements $\text{deg}(a_{ij}) \leq d$ we can always construct a matrix C , which defines an equivalent system, such that*

$$CA = \begin{pmatrix} C_1 & 0 \\ C_2 & E \end{pmatrix} A = \left(\begin{array}{c|c} a_1 & 0 \\ \vdots & \star \\ 0 & a_r \\ \hline 0 & 0 \end{array} \right) \tag{1}$$

where E is the identity matrix.

Due to this lemma, we can assume that our system is equivalent to a system in diagonal form:

$$a_k V_k + \sum_{r+1 \leq l \leq s} a_{k,l} V_l = b_k, \quad 1 \leq k \leq r, \quad \deg(a_k), \deg(a_{k,l}), \deg(b_k) \leq 2pmd.$$

4.2 Normalization

Once the system is in diagonal form, we need to normalize it. To do this, we construct some syzygies, applying Lemma 1 to the submatrix of the first r columns and the column $l > r$. There always exist $h^{(l)}, h_1^{(l)}, \dots, h_r^{(l)}$ such that:

$$a_k h_k^{(l)} + a_{k,l} h^{(l)} = 0, \quad 1 \leq k \leq r \quad \deg(h^{(l)}), \deg(h_i^{(l)}) \leq 4p^2 m^2 d$$

The result that gives the normalization in the Weyl algebra is the following one:

Lemma 4 ([Gr1], Lemma 4). *Given $g_1, \dots, g_t \in D_n$ a family of elements, there is a nonsingular linear transformation of $2n$ -dimensional space with basis $x_1, \dots, x_n, \partial_1, \dots, \partial_n$ under which:*

$$\begin{aligned} x_i &\rightarrow \Gamma_{x_i} = \sum_{j=1}^n \gamma_{i,j}^{(1,1)} x_j + \sum_{j=1}^n \gamma_{i,j}^{(1,2)} \partial_j; \\ \partial_i &\rightarrow \Gamma_{\partial_i} = \sum_{j=1}^n \gamma_{i,j}^{(2,1)} x_j + \sum_{j=1}^n \gamma_{i,j}^{(2,2)} \partial_j \end{aligned}$$

such that the following relations hold:

$$\begin{aligned} \Gamma_{x_i} \Gamma_{\partial_i} &= \Gamma_{\partial_i} \Gamma_{x_i} - 1; & \Gamma_{x_i} \Gamma_{x_j} &= \Gamma_{x_j} \Gamma_{x_i} \\ \Gamma_{\partial_i} \Gamma_{\partial_j} &= \Gamma_{\partial_j} \Gamma_{\partial_i} & \Gamma_{\partial_i} \Gamma_{x_j} &= \Gamma_{x_j} \Gamma_{\partial_i} \quad i \neq j, \end{aligned}$$

and if we denote by Γ_{g_i} the transformed of g_i with the indicated linear transformation, we have $\Gamma_{g_i} = \partial_n^{\deg(g_i)} + \widetilde{\Gamma}_{g_i}$.

Remark 1. The main fact in the proof of the last Lemma 4 is that the matrices of the linear transformations defined by the relations in the Weyl algebra are a transitive group.

Let $\{g_1, \dots, g_t\}$ be a set of elements in $R = \mathbf{C}[s, t, x_1, \dots, x_n, \partial_1, \dots, \partial_n]$. Let us see why we can not assure the existence of a linear transformation Γ that produces

$$\Gamma_{g_i} = v^{\deg(g_i)} + \widetilde{\Gamma}_{g_i},$$

where v is a single variable.

A general linear transformation as the one postulated in Lemma 4 has the form:

$$\begin{aligned} s &\rightarrow \Gamma_s = \alpha_1 s + \beta_1 t + \sum_{j=1}^n \gamma_j^{(s,1)} x_j + \sum_{j=1}^n \gamma_j^{(s,2)} \partial_j \\ t &\rightarrow \Gamma_t = \alpha_2 s + \beta_2 t + \sum_{j=1}^n \gamma_j^{(t,1)} x_j + \sum_{j=1}^n \gamma_j^{(t,2)} \partial_j \\ x_i &\rightarrow \Gamma_{x_i} = \alpha_i^{(1)} s + \beta_i^{(1)} t + \sum_{j=1}^n \gamma_{i,j}^{(1,1)} x_j + \sum_{j=1}^n \gamma_{i,j}^{(1,2)} \partial_j \\ \partial_i &\rightarrow \Gamma_{\partial_i} = \alpha_i^{(2)} s + \beta_i^{(2)} t + \sum_{j=1}^n \gamma_{i,j}^{(2,1)} x_j + \sum_{j=1}^n \gamma_{i,j}^{(2,2)} \partial_j \end{aligned}$$

and it has to verify the following relations:

- (1) $\Gamma_s \Gamma_t = \Gamma_t \Gamma_s + \Gamma_t$; (2) $\Gamma_s \Gamma_{x_i} = \Gamma_{x_i} \Gamma_s$; (3) $\Gamma_s \Gamma_{\partial_i} = \Gamma_{\partial_i} \Gamma_s$;
- (4) $\Gamma_t \Gamma_{x_i} = \Gamma_{x_i} \Gamma_t$; (5) $\Gamma_t \Gamma_{\partial_i} = \Gamma_{\partial_i} \Gamma_t$; (6) $\Gamma_{x_i} \Gamma_{\partial_i} = \Gamma_{\partial_i} \Gamma_{x_i} - 1$;
- (7) $\Gamma_{x_i} \Gamma_{x_j} = \Gamma_{x_j} \Gamma_{x_i}$; (8) $\Gamma_{\partial_i} \Gamma_{\partial_j} = \Gamma_{\partial_j} \Gamma_{\partial_i}$; (9) $\Gamma_{x_i} \Gamma_{\partial_j} = \Gamma_{\partial_j} \Gamma_{x_i}$

From relation (1), we obtain $\alpha_2 = \gamma_j^{(t,1)} = \gamma_j^{(t,2)} = 0$ for all j , so $\Gamma_t = \beta_2 t$. The change must be nonsingular, so we have $\beta_2 \neq 0$, and again using relation (1) we deduce that $\alpha_1 = 1$. Using relation (4), we obtain that $\alpha_i^{(1)} = 0$ for all i , and with relation (5) that $\alpha_i^{(2)} = 0$ for all i .

By relation (2) (Γ_s commutes with Γ_{x_i}) we have $\beta_i^{(1)} = 0$, and relation (3) gives $\beta_i^{(2)} = 0$. So Γ must verify:

$$\begin{aligned} s &\rightarrow \Gamma_s = s + \beta_1 t + \sum_{j=1}^n \gamma_j^{(s,1)} x_j + \sum_{j=1}^n \gamma_j^{(s,2)} \partial_j \\ t &\rightarrow \Gamma_t = \beta_2 t \\ x_i &\rightarrow \Gamma_{x_i} = \sum_{j=1}^n \gamma_{i,j}^{(1,1)} x_j + \sum_{j=1}^n \gamma_{i,j}^{(1,2)} \partial_j \\ \partial_i &\rightarrow \Gamma_{\partial_i} = \sum_{j=1}^n \gamma_{i,j}^{(2,1)} x_j + \sum_{j=1}^n \gamma_{i,j}^{(2,2)} \partial_j \end{aligned}$$

Due to relations from (6) to (9) (between Γ_{x_i} and Γ_{∂_j}) we have that the submatrix

$$\begin{pmatrix} \gamma_{i,j}^{(1,1)} & \gamma_{i,j}^{(1,2)} \\ \gamma_{i,j}^{(2,1)} & \gamma_{i,j}^{(2,2)} \end{pmatrix}$$

verifies the relations of Lemma 4, and in addition, from the relations with Γ_s it verifies

$$\sum \gamma_i^{(s,1)} \gamma_{i,i}^{(1,2)} = \sum \gamma_i^{(s,2)} \gamma_{i,i}^{(1,1)} \quad \sum \gamma_i^{(s,1)} \gamma_{i,i}^{(2,2)} = \sum \gamma_i^{(s,2)} \gamma_{i,i}^{(2,1)}.$$

Anyway if we take for example tx_1 , the requirements for Γ produce

$$\Gamma_{tx_1} = \beta_2 t \Gamma_{x_1} \neq v^2 + \widetilde{\Gamma}_{tx_1}.$$

Thus we can not repeat the second step of the process in our PBW algebra in the same way that appears in [Gr1].

Problem 1. Find a general bound for the solutions of a general linear system over any PBW algebra or, at least, give such a bound for R .

We will not treat this general problem: with the aim of obtaining a bound for the complexity of the annihilating ideal of f^s , we will consider only the particular case of one equation of the type that would produce the definition of the ideal I in section 2.1 or I' in section 2.2. In both cases we are interested in the complexity of computing their Gröbner bases (in different rings), and we do it considering the equivalent problem of computing the *syzygies* of the generators of our respective ideals.

Note 1. In the algorithm of [OT1] the calculations are computed in a Weyl algebra of $2n + 4p$ variables in all, or more precisely in a commutative polynomial ring with $n + 3p$, (x, u, v, t) commutative variables extended with $n + p$, (∂_x, ∂_t) “differential” variables. Let us denote by A this algebra. The complexity of computing the annihilating ideal of f^s is bounded by the complexity of computing a Gröbner basis in A .

Recall that the complexity in the Weyl algebra is given by the following theorem:

Theorem 3 (Th. 6,[Gr1]). *Given a solvable system in the Weyl algebra D_n :*

$$\sum_{1 \leq l \leq s} u_{k,l} V_l = w_k, \quad 1 \leq k \leq m$$

with $\deg(u_{k,l}), \deg(w_k) \leq d$. *There exists a solution with $\deg(V_i) < (md)^{2^{O(n)}}$*

As we said before in the Briançon-Maisonobe ring R we can not construct a similar algorithm to bound the degree of a solution for a system in general. But in our very special case, our problem is equivalent to computing the solutions of the equation:

$$(s_1 + f_1 t_1) V_1 + \dots + (s_p + f_p t_p) V_p + (\partial_1 + \sum_j \frac{\partial f_j}{\partial x_1} t_j) V_{p+1} + \dots + (\partial_n + \sum_j \frac{\partial f_j}{\partial x_n} t_j) V_{p+n} = 0$$

To simplify notation we write the precedent equation as

$$\sum_l Q_l V_l = 0 \tag{2}$$

Theorem 4. *Given $f = (f_1, \dots, f_p)$, the complexity of the computation of the annihilating ideal of f^s in the Briançon-Maisonobe algebra $R = D_n[s_1, \dots, s_p, t_1, \dots, t_p]$ is bounded by the complexity of the computation of the syzygies of the elements $\partial_i + \sum_j \frac{\partial f_j}{\partial x_i} t_j$ in the Weyl algebra $D_n[t_1, \dots, t_p]$.*

Proof. We follow the notations of [Gr1] in this proof. We first compute $h_1^{(l)}, h^{(l)}$ for $2 \leq l \leq n + p$ such that:

$$\begin{aligned} (s_1 + f_1 t_1)h_1^{(2)} + (s_2 + f_2 t_2)h^{(2)} &= 0 \\ &\vdots \\ (s_1 + f_1 t_1)h_1^{(p)} + (s_p + f_p t_p)h^{(p)} &= 0 \\ (s_1 + f_1 t_1)h_1^{(p+1)} + (\partial_1 + \sum_j \frac{\partial f_j}{\partial x_1} t_j)h^{(p+1)} &= 0 \\ &\vdots \\ (s_1 + f_1 t_1)h_1^{(p+n)} + (\partial_n + \sum_j \frac{\partial f_j}{\partial x_n} t_j)h^{(p+n)} &= 0. \end{aligned}$$

The aim of these $h^{(l)}$ is to reduce any solution $V = (V_1, \dots, V_{p+n})$ of equation (2) to another one without s_1 from which you can recover V . The process will be repeated for s_2, \dots, s_p .

It is easy to see that

$$[s_i + f_i t_i, s_j + f_j t_j] = 0$$

$$\begin{aligned} [s_i + f_i t_i, \partial_j + \sum_l \frac{\partial f_l}{\partial x_j} t_l] &= s_i (\sum_l \frac{\partial f_l}{\partial x_j} t_l) + f_i t_i \partial_j - \partial_j f_i t_i - (\sum_l \frac{\partial f_l}{\partial x_j} t_l) s_i = \\ &= t_i s_i \frac{\partial f_i}{\partial x_j} + t_i \frac{\partial f_i}{\partial x_j} + \sum_{l \neq i} t_l s_i \frac{\partial f_l}{\partial x_j} + t_i f_i \partial_j - t_i f_i \partial_j - t_i \frac{\partial f_i}{\partial x_j} - \sum_l \frac{\partial f_l}{\partial x_j} t_l s_i = 0. \end{aligned}$$

Let us define $h^{(l)} = s_1 + f_1 t_1$ for all $l \geq 2$. We make the division of the V_l of equation (2), $l \geq 2$ by $h^{(l)}$ with respect to a lexicographical ordering with s_1 greater than any other variable. We obtain a remainder \bar{V}_l such that $\text{deg}_{s_1}(\bar{V}_l) < \text{deg}_{s_1}(h^{(l)}) = 1$, so it has no s_1 . So $V_l = h^{(l)} \bar{\bar{V}}_l + \bar{V}_l$, and adding the relation $Q_1 h_1^{(l)} + Q_l h^{(l)} = 0$ multiplied by $-\bar{\bar{V}}_l$ to equation (2), we obtain:

$$Q_1 \bar{V}_1 + Q_2 \bar{V}_2 + \dots + Q_{n+p} \bar{V}_{n+p} = 0$$

with Q_i, \bar{V}_i without s_1 for $i \geq 2$, so $\bar{V}_1 = 0$, where $\bar{V}_1 = V_1 - h_1^{(2)} \bar{\bar{V}}_2 - \dots - h_1^{(n+p)} \bar{\bar{V}}_{n+p}$, and we have the new equation:

$$Q_2 \bar{V}_2 + \dots + Q_{n+p} \bar{V}_{n+p} = 0$$

in a Briançon-Maisonobe algebra $\mathbf{C}[s_2, \dots, s_p, t_1, \dots, t_p, x, \partial]$.

Repeating the process for Q_2, \dots, Q_p , we reduce our problem to solving:

$$(\partial_1 + \sum_j \frac{\partial f_j}{\partial x_1} t_j) V_{p+1} + \dots + (\partial_n + \sum_j \frac{\partial f_j}{\partial x_n} t_j) V_{p+n} = 0$$

in the Weyl algebra $D_n[t_1, \dots, t_p]$.

As a consequence of 4, the bound for the complexity of computing the annihilating ideal of f^s in R is bounded by the complexity of computing a Gröbner basis in a Weyl algebra with $3p$ variables less than the one required by the method of [OT1]. Although the complexity of computing these objects in any case is known to be double exponential (with respect to the number of variables and the total degree of the generators of the ideal) by Theorem 3, it is clear that the reduction of $3p$ variables of [BM1] is a significant advantage in practice as it is shown in the examples (see section 3). The theoretical superiority of the method of [BM1] is an open problem.

Problem 2. Is the bound proposed in this work is reached *a la* Mayr-Meyer ([MM1])? (that is to say, find an example of annihilating ideal with this worst complexity).

References

- [Ba1] Bahloul, R. Algorithm for computing Bernstein-Sato ideals associated with a polynomial mapping. *Journal of Symbolic Computation* **32**, 643–662, 2001.
- [Be1] Bernstein, I.N. The analytic continuation of generalized functions with respect to a parameter. *Funct. Anal.* **6** 273–285, 1972.
- [Bj1] Bjork, J.E. *Dimensions over Algebras of Differential Operators*. Preprint, 1973.
- [BM1] Briançon, J. and Maisonobe, Ph. *Remarques sur l'idéal de Bernstein associé à des polynômes*. PUMA, 650 (preprint). 2002.
- [BS1] Budur, N. and Saito M. Multiplier ideals, V -filtration and spectrum. arXiv:math.AG/0305118, 2003.
- [BGV1] Bueso, J.L. Gómez-Torrecillas, J. and Verschoren, A. *Algorithmic methods in non-commutative algebra. Applications to quantum groups*. Mathematical modelling. Theory and applications **17**, Kluwer Academic Publishers. 2004.
- [GHU1] Gago-Vargas, J., Hartillo-Hermoso, M.I. and Ucha-Enríquez, J.M. On the computation of the annihilating ideal of a product of polynomials. In preparation.
- [GLS1] G.-M. Greuel, V. Levandovskyy, and H. Schönemann. SINGULAR::PLURAL 2.1. A Computer Algebra System for Noncommutative Polynomial Algebras. Centre for Computer Algebra, University of Kaiserslautern (2003). <http://www.singular.uni-kl.de/plural>.
- [Gr1] Grigoriev, D. Complexity of solving systems of linear equations over the rings of differential operators. In *Effective methods in algebraic geometry (Castiglione, 1990)*, Progr. Math., **94**, Birkhauser Boston, MA 195–202. 1991.
- [H1] Hamm, H.A. Remarks on asymptotic integrals, the polynomial of I.N. Bernstein and the Picard-Lefschetz monodromy. In *Several complex variables (Proc. Sympos. Pure Math. Vol XXX, Part 1, Williams Coll., Williamstown, Mass. 1975)*, 31–35. Amer. Math. Soc., Providence, R.I. 1977.
- [K1] Kashiwara, M. B-functions and holonomic systems. Rationality of roots of B-functions. *Invent. Math.* **38** (1) 33–53. 1976.
- [L1] Lichtin, B. Generalized Dirichlet series and b-functions. *Compositio Math* **65** 81–120. 1988.

- [Mal1] Malgrange, B. Le polynôme de Bernstein d'une singularité isolée. In *Fourier integral operators and partial differential equations (Colloq. Internat., Univ. Nice, Nice, 1974)*. Lecture Notes in Math. **459** Springer-Verlag, New York 98–119. 1975.
- [MM1] Mayr, E.W. and Meyer, A.R. The complexity of the word problems for commutative semigroups and polynomial ideals. *Adv. in Math.* **46(3)**, 305–329. 1982.
- [N1] M. Noro, T. Shimoyama and Takeshima, T. A Computer Algebra System **Risa/Asir** (2000). Available at <ftp://archives.cs.ehime-u.ac.jp/pub/asir2000/>.
- [O1] Oaku, T. An algorithm of computing b-functions. *Duke Math. J.* **87** 115–132 1997.
- [OT1] Oaku, T. Takayama, N. An algorithm for de Rham cohomology groups of the complement of an affine variety via D-module computation. *Journal of Pure and Applied Algebra* **139** 201–233 1999.
- [Sa1] Sabbah, C. Proximité évanescence I. La structure polaire d'un D-Module. Appendice en collaboration avec F.J. Castro Jiménez. *Compositio Math.* **62**, 283–328. 1987.
- [Sa2] Sabbah, C. Proximité évanescence II. Equations fonctionnelles pour plusieurs fonctions analytiques. *Compositio Math.* **64**, 213–241. 1987.
- [Se1] Seidenberg, A. Constructions in algebra. *Trans. Amer. Math. Soc.* **197** 273–313. 1974.
- [St1] Stafford, J. T. Module structure of Weyl algebras. *J. London Math. Soc.* **(2) 18**, no. 3, 429–442. 1978.