

Note

## Counterexample to a conjecture of Györi on $C_{2l}$ -free bipartite graphs<sup>☆</sup>

C. Balbuena<sup>a</sup>, P. García-Vázquez<sup>b</sup>, X. Marcote<sup>a</sup>, J.C. Valenzuela<sup>c</sup>

<sup>a</sup>Departament de Matemàtica Aplicada III, Universitat Politècnica de Catalunya, Campus Nord, Edifici C2, C/ Jordi Girona 1 i 3, E-08034 Barcelona, Spain

<sup>b</sup>Departamento de Matemática Aplicada I, Universidad de Sevilla, Avda Reina Mercedes 2, E-41012 Sevilla, Spain

<sup>c</sup>Departamento de Matemáticas, Universidad de Cádiz, Avda Ramón Puyol s/n, E-11202 Cádiz, Spain

Received 24 November 2005; received in revised form 5 July 2006; accepted 10 July 2006  
Available online 24 August 2006

### Abstract

A counterexample on a conjecture of Györi related with  $C_{2l}$ -free bipartite graphs is described.  
© 2006 Elsevier B.V. All rights reserved.

*Keywords:* Cycles in bipartite graphs; Forbidden subgraphs

In 1997 Györi [2] studied the structure of  $C_6$ -free bipartite graphs and the relationship between this problem and some interesting results of Erdős et al. [1] on a number-theoretic problem. Namely, Györi proved a conjecture of Erdős et al. [1] regarding the maximum number of edges that a  $C_6$ -free bipartite graph can have. Moreover, he proved another theorem that generalizes the previous one for cycles of longer length. In this paper Györi stated a conjecture [2, p. 373] that apparently contains a misprint<sup>1</sup> and it should have been expressed in this way:

**Conjecture 1.** If  $G = (X, Y)$  is a bipartite graph with color classes  $X, Y$  where  $|X| = m$ ,  $|Y| = n$ ,  $m^2 \leq n$ ,  $3 \leq l \leq m$  and  $G$  has at least  $(l - 1)n + m - l + 2$  edges, then  $G$  must contain a cycle of length  $2l$ .

In a recent paper [3], the same author disproves Conjecture 1 for  $l = 3$ , but leaves the proof or refutation for  $l \geq 4$  as an open problem. In this note we provide a counterexample that disproves Conjecture 1 when  $m \geq 2l - 1$ . Let us denote by  $K_{(m,n)}$  the complete bipartite graph with  $m$  vertices in the first class and  $n$  vertices in the second one. Let us also denote by  $d_G(v)$  the degree of the vertex  $v$  in the graph  $G$ .

Let  $l, m$  be integers such that  $3 \leq l \leq 2(l - 1) \leq m$ . We consider the graphs  $G_1 = K_{(m-l+1, l-1)}$  and  $G_2 = K_{(l-1, n-l+2)}$ . Take two vertices  $u \in V(G_1)$  and  $v \in V(G_2)$  with  $d_{G_1}(u) = m - l + 1$  and  $d_{G_2}(v) = l - 1$ , and let  $G$  be the bipartite

<sup>☆</sup> Research supported by the Ministry of Education and Science, Spain, and the European Regional Development Fund (ERDF) under project MTM2005-08990-C02-02.

*E-mail addresses:* [m.camino.balbuena@upc.edu](mailto:m.camino.balbuena@upc.edu) (C. Balbuena), [pgvazquez@us.es](mailto:pgvazquez@us.es) (P. García-Vázquez), [francisco.javier.marcote@upc.edu](mailto:francisco.javier.marcote@upc.edu) (X. Marcote), [jcarlos.valenzuela@uca.es](mailto:jcarlos.valenzuela@uca.es) (J.C. Valenzuela).

<sup>1</sup> The original conjectured value  $(l - 1)n + m - l + 1$  (see [2]) may be easily disproved by means of a graph roughly outlined by the author. We appreciate the referee's comments enlightening this misprint.




Fig. 1. The graph  $G$  with  $l = 3$ ,  $m = 5$  and  $n = 25$ .

graph on  $m$  and  $n$  vertices obtained by gluing the graphs  $G_1$  and  $G_2$  in such a way that the vertex  $u$  of  $G_1$  is identified with the vertex  $v$  of  $G_2$  (see Fig. 1).

Clearly, any cycle of  $G$  must be entirely contained in either  $G_1$  or  $G_2$ . But  $G_i$ ,  $i = 1, 2$ , cannot contain a cycle of length  $2l$  because one of its classes has cardinality  $l - 1$ . So  $G$  is free of  $C_{2l}$  and it has size  $e(G) = (m - l + 1)(l - 1) + (l - 1)(n - l + 2) \geq (l - 1)n + m - l + 2$  because  $m \geq 2l - 1$ . Therefore, Conjecture 1 is disproved for  $m \geq 2l - 1$ .

In [2], Györi proved the following result:

**Theorem.** *If  $G(X, Y)$  is a bipartite graph with color classes  $X, Y$  such that  $|X| = m$ ,  $|Y| = n$ ,  $m^2 \leq n$  and  $G$  has at least  $(l - 1)n + c(l)m^2$  edges for some constant  $c(l)$  then  $G$  must contain a cycle of length  $2l$ .*

Thus, we propose the following reformulation of the conjecture:

**Conjecture 2.** *If  $G = (X, Y)$  is a bipartite graph with color classes  $X, Y$  where  $|X| = m$ ,  $|Y| = n$ ,  $m^2 \leq n$ ,  $3 \leq l \leq m$  such that  $m > (l - 1)^2$  and  $G$  has at least  $(l - 1)n + 1/(l - 1)m^2$  edges then  $G$  must contain a cycle of length  $2l$ .*

Corollary of Theorem 1 in [3] confirms our Conjecture 2 for  $l = 3$ . For  $l \geq 4$  it is still an open problem.

**References**

[1] P. Erdős, A. Sarközy, V.T. Sós, On product representation of powers I, European J. Combin. 16 (1995) 567–588.  
 [2] E. Györi,  $C_6$ -free bipartite graphs and product representation of squares, Discrete Math. 165/166 (1997) 371–375.  
 [3] E. Györi, Triangle-free hypergraphs, Combinatorics, Probab. Comput. 15 (1–2) (2006) 185–191.